

Testing the Matter Bounce with Primordial Non-Gaussianity: A SPHEREx Sensitivity Recast with a MegaMapper Outlook

Houston Golden^{1, *}

¹*Independent Researcher, Los Angeles, California, USA*

(Dated: July 6, 2026)

Scope. This work is a sensitivity recast of a single externally published forecast, not an independent forecast: every quoted SPHEREx significance and Bayes factor rescales the imported Heinrich *et al.* [1] $\sigma(f_{\text{NL}}^{\text{local}}) \approx 0.7$ baseline by the explicit template-mismatch recast $(\hat{f}_{\text{NL}}^{\text{bounce}}, \sigma^{\text{bounce}}) = (\hat{f}_{\text{NL}}^{\text{local}}/r, \sigma^{\text{local}}/r)$ with $r = 0.84$ (Eq. 5) plus a heuristic additive-quadrature systematic budget, with no independent bispectrum Fisher matrix constructed here and cross-parameter correlations neglected in the recast. The headline ranges are therefore conditional sensitivity envelopes, not internally derived measurement precisions; because they are drawn from qualitatively different null procedures (signal-only CMB-Fisher weighting, LSS noise-weighting, and b_ϕ /GR-marginalized budgets), the individual σ endpoints within a range are *not directly comparable* to one another and should be read as a scoping envelope. *Load-bearing caveat* (\star): the entire forecast is conditional on assumption (d) — faithful third-order (cubic) bispectrum transmission through the bounce — which is verified at linear order [2] and, at cubic order, now *derived* to a bounded systematic: effective LQC adds no new scalar degree of freedom [2, 3], so the Wilson-Ewing bounce is single-clock, nonlinear superhorizon ζ -conservation (Lyth-Malik-Sasaki / Maldacena- δN) holds to all orders, and transmission = $1 \pm O((k\eta_{\text{bounce}})^2) \approx 1 \pm 10^{-4}$ ($\delta f_{\text{NL}} \lesssim 10^{-3}$, negligible against $\sigma(f_{\text{NL}}) \approx 0.7$); this model also evades the Quintin *et al.* [4] bounce no-go. The one remaining model-dependence is the sign of the subleading gradient coefficient (a citable quantization choice, Lorentzian $c_s^2 = 1$ in the dressed-metric scheme adopted here), not an open computation. With that framing fixed: a matter-dominated contracting phase preceding a nonsingular bounce produces a minimally parameterized local-type non-Gaussianity $f_{\text{NL}}^{\text{local}} = -35/16 = -2.1875$, in the scalar-only matter-bounce class defined by assumptions (a)–(f) of Sec. II C — in particular (e) no prolonged post-bounce inflation, (f) negligible fermion-sourced torsion during contraction, and (d) faithful third-order bispectrum transmission through the bounce (verified at linear order [2] and supported at cubic order by a superhorizon-scaling estimate). The bounce-vs-inflation discrimination is dual-pronged: in the Planck/local-template gauge convention used by SPHEREx and BOSS, the slow-roll value $f_{\text{NL}}^{\text{inf}} \approx 0.015$ at $n_s = 0.9649$ [5, 6] gives a contrast $|f_{\text{NL}}^{\text{bounce}}|/|f_{\text{NL}}^{\text{inf}}| \approx 146$; in the conformal-Fermi physical-observer frame the squeezed-limit consistency relation [7, 8] sends $f_{\text{NL}}^{\text{local}} \rightarrow 0$ for single-field slow-roll, so a non-zero physical-frame local detection (modulo CFC gradient/projection corrections) would disfavor the single-field slow-roll attractor. The forecast estimators measure the gauge-frame f_{NL} directly; the CFC statement is a complementary theoretical discriminator, not the on-sky observable.

We forecast tests of this prediction with SPHEREx (launched March 2025, primary survey through ~ 2027 , first PNG-suitable release expected ~ 2028) and the proposed MegaMapper via scale-dependent bias and the galaxy bispectrum. A long-standing factor-of-two discrepancy in the literature — Cai *et al.* [9] report $f_{\text{NL}}^{\text{local}} = -35/8$ while Li *et al.* [10] report $-35/16$ (at $c_s = 1$), both in the identical $f_{\text{NL}} = 10A/(3\sum k_i^3)$ normalization and the identical squeezed limit — *is resolved here in favour of $-35/16$* , which we accordingly adopt as the central matter-bounce value. Re-summing Cai *et al.*'s own four cubic-action vertex contributions (field redefinition, $\zeta\dot{\zeta}^2$, $\dot{\zeta}\partial\zeta\partial\chi$, $\zeta(\partial_i\partial_j\chi)^2$) at $\epsilon = 3/2$ and taking the squeezed limit $k_1 \ll k_2 = k_3$ yields a clean $f_{\text{NL}}^{\text{local}} = -35/16$ (with an $O(k_1^2/k^2)$ correction), in exact agreement with the general- c_s formula of Li *et al.* [10] (their Eq. 5.1, $-165/16 + 65/(8c_s^2) \rightarrow -35/16$ at $c_s = 1$); the published $-35/8$ arises only from a spurious $+(99/128)\sum_i k_i^3$ local-shaped term introduced when Cai *et al.* collapse their (correct) ϵ -order-grouped expressions into a single final polynomial (their Eq. 37), a term absent from every intermediate expression in that same paper. Both works use the identical $f_{\text{NL}} = 10A/(3\sum k_i^3)$ convention, so this is an arithmetic error, not a convention difference (Appendix A). This settles the 8-year discrepancy and *halves* the headline matter-bounce amplitude relative to the erroneous $-35/8$. We quantify the template mismatch between the matter-bounce and local templates: a local estimator recovers 83%–88% of the bounce signal across noise-weighting schemes ($r \in [0.829, 0.876]$); CMB Fisher signal-only $r = 0.876$, LSS/SPHEREx noise-weighted $r \approx 0.83$), validated via ℓ -space Fisher overlap, 200 injection-recovery realizations, and a 10,000-sample null-space scan of the underdetermined polynomial coefficients (shape cosine $r_{\text{cos}} > 0.97$ for all samples). Only the noise-weighted $r \approx 0.83$ enters the SPHEREx significance; r is applied as a shape-weighted degradation to the Heinrich *et al.* baseline rather than recomputed as an independent cross-Fisher matrix, making this a sensitivity recast rather than an independent forecast.

The Heinrich *et al.* multi-tracer galaxy bispectrum forecast under the local-template normalization $B_\zeta^{\text{local}}(k_1, k_2, k_3) = (6f_{\text{NL}}^{\text{local}}/5)[P_\zeta(k_1)P_\zeta(k_2) + 2 \text{ perms}]$ achieves $\sigma(f_{\text{NL}}^{\text{local}}) \approx 0.7$ [1]. After

template-mismatch correction we obtain bispectrum-only $\sim 2.6\text{--}2.75\sigma$ at the central $f_{\text{NL}} = -35/16$ (LSS noise-weighted to CMB Fisher endpoints; $|-35/16|r/\sigma$ with $r = 0.84\text{--}0.876$ and $\sigma(f_{\text{NL}}) = 0.7$), reducing to a realistic $\sim 1.3\text{--}2.75\sigma$ after the systematic budget (mismatch, ϵ -correction, polynomial-null-space scatter ± 0.13 in r at fixed sampling measure with 16th-percentile $r_{16\text{th}} = 0.70$ as a distributional robustness bound separate from the conservative floor which uses the noise-weighted central $r = 0.84$ of Table IV, photometric- z degradation, PNG-bias b_ϕ marginalization, and relativistic projection); these systematics are combined additively in quadrature, assuming the individual contributions are uncorrelated, so the realistic $1.3\text{--}2.75\sigma$ range is a scoping sensitivity envelope under an additive-quadrature heuristic systematic budget, *not* a joint-covariance forecasted measurement precision (the true joint constraint could be weaker, or marginally stronger, than this envelope); a full bispectrum joint Fisher over the systematic nuisances is not performed here. Moreover, the imported Heinrich *et al.* multi-tracer covariance was constructed for a purely local template; any additional variance from the non-local tails of the bounce shape in the full multi-tracer estimator covariance is not modeled and is absorbed into this systematic envelope (a full re-derivation of the Heinrich Fisher at the bounce fiducial, which would convert this into a derived bound, is out of scope and flagged for follow-up). Separately, and as an independently computed result kept distinct from this bispectrum-only headline, a joint scale-dependent-bias (SDB) Fisher matrix for $(f_{\text{NL}}, n_{f_{\text{NL}}})$ (`c8.fn1.running.fisher.json`, Planck 2018, CAMB 1.6.6; SDB signal of §III, forecast reported in §IX) shows that the $f_{\text{NL}}\text{--}n_{f_{\text{NL}}}$ running degeneracy — and, with the per-sample linear biases co-marginalized, the $b_1\text{--}f_{\text{NL}}$ degeneracy — *loosen* the SDB f_{NL} constraint by $2.0\text{--}4.6\times$ ($\sigma_{\text{unmarg}}(f_{\text{NL}}) = 1.53 \rightarrow \sigma_{\text{marg}}(f_{\text{NL}}) = 3.08$ fixed-bias $\rightarrow 7.06$ bias-marginalized; anti-correlation $\rho \approx -0.87$ to -0.97), with the running the dominant direction. This computed SDB result is a subordinate cross-check on the scale dependence, not a competitor to the bispectrum-only headline, and the bispectrum template-overlap factor $r = 0.84$ does not apply to it. With the Cai-Li factor-of-two now resolved in favour of $f_{\text{NL}}^{\text{local}} = -35/16$ (Appendix A), we headline a single central amplitude rather than a dual-value range: the bispectrum-only significance is $\sim 2.6\text{--}2.75\sigma$ (LSS noise-weighted to CMB Fisher endpoints) and the realistic post-systematic-budget significance is $\sim 1.3\text{--}2.75\sigma$; the erroneous $-35/8$ value (which would have doubled these to $5.2\text{--}5.5\sigma / 2.6\text{--}5.5\sigma$) is retained only as a footnote/upper-bookkeeping reference in Appendix A, not as a forecast input. Both endpoints rest on the single imported Heinrich *et al.* $\sigma(f_{\text{NL}}^{\text{local}}) \approx 0.7$ baseline [1] recast for template mismatch, and are a sensitivity recast rather than an independent cross-Fisher forecast. MegaMapper, proposed but not yet funded, could reach $\sigma(f_{\text{NL}}) \approx 0.5$ ideally, projecting an illustrative $1.5\text{--}3.5\sigma$ envelope that reflects design uncertainty as much as measurement uncertainty.

A closed-form Bayesian comparison validated across three independent 10^5 -realization Monte Carlo ensembles (§VI, Table II) finds that a SPHEREx detection near the central prediction $f_{\text{NL}} = -35/16 = -2.1875$ favors the bounce over tuned multifield competitors at Bayes factor $\text{BF} \approx 9$ (recommended $\sigma_{\text{theory}} = 1.0$ Gaussian bounce prior, broad multifield $[-15, +15]$ competitor) up to $\text{BF} \approx 14$ at the delta-prior theoretical maximum, evaluated under the noise-weighted $r \approx 0.84$ template-mismatch bookkeeping consistent with the significance headline (the $r \rightarrow 1$ bookkeeping endpoint gives $\text{BF} \approx 10\text{--}17$ and is reported in Table II); Table II reports the $r \rightarrow 1$ endpoint values while the abstract headline applies the noise-weighted $r \approx 0.84$ rebooking $\sigma_{\text{eff}} = \sigma(f_{\text{NL}}^{\text{local}})/r$ to those entries; a curvaton-natural $[-5, +5]$ competitor narrows this to $\text{BF} \approx 4\text{--}7$. These Bayes factors should be read as illustrative of the discriminating power available given the current theoretical uncertainty in the bounce prediction — not as definitive model-selection evidence — and as $\gg 1$ against the parameter-free single-field slow-roll prediction. A SPHEREx null would disfavor the quasi-dust matter bounce benchmark at the same post-systematic-budget level as a detection ($\sim 1.3\text{--}2.75\sigma$ at the central $f_{\text{NL}} = -35/16$; the exclusion arithmetic is symmetric).

The central value $f_{\text{NL}}^{\text{local}} = -35/16$ is the resolution of the historical Cai-Li factor-of-two: a from-scratch re-summation of Cai *et al.*'s [9] own four cubic-action vertices, plus the independent general- c_s derivation of Li *et al.* [10], both give $-35/16$, while the published $-35/8$ is traced to a single spurious $+(99/128)\sum_i k_i^3$ term in Cai's final combined polynomial (Appendix A). Adopting the corrected $-35/16$ gives a matter-bounce SPHEREx significance of $\sim 2.6\text{--}2.75\sigma$ optimistic (bispectrum-only, template-corrected) and $\sim 1.3\text{--}2.75\sigma$ after the full systematic budget — preserving qualitative discriminatory power against single-field inflation ($|f_{\text{NL}}^{\text{inf}}| \approx 0.015$, opposite sign) while dropping the erroneous $\geq 5\sigma$ headline that the $-35/8$ value would have produced. The corrected amplitude therefore sets a real, not merely hypothetical, sensitivity target for SPHEREx (see Appendix A and Sec. II C).

I. INTRODUCTION

The inflationary paradigm provides a remarkably successful framework for generating the observed spec-

* houston@hubify.com

trum of primordial perturbations. Standard single-field slow-roll inflation predicts a nearly scale-invariant, nearly Gaussian spectrum with a small, positive local-type non-Gaussianity $f_{\text{NL}} \approx (5/12)(1 - n_s) \approx 0.015$, set by the Maldacena consistency relation [5] (gauge-frame value; the conformal-Fermi-frame equivalent differs by $\mathcal{O}(\text{slow-roll})$ corrections [7, 8] and is not the bounce-discriminating quantity here, since the bounce-vs-inflation contrast remains $|f_{\text{NL}}^{\text{bounce}}| \gg |f_{\text{NL}}^{\text{inf}}|$ in either frame).

Bouncing cosmology offers an alternative origin for primordial perturbations: modes exit the Hubble radius during a contracting phase and re-enter after a nonsingular bounce. In particular, a matter-dominated contraction ($w \approx 0$) produces a scale-invariant scalar spectrum through the growth of the curvature perturbation ζ on superhorizon scales [11, 12].

A distinctive prediction of the matter bounce is a large, negative, minimally parameterized local-type non-Gaussianity $f_{\text{NL}} = -35/16 = -2.1875$ [9, 10, 13] at zeroth order in the ϵ -expansion (the corrected value; the earlier literature $-35/8$ contains a factor-of-two arithmetic error, resolved in Appendix A) (the first-order coefficient κ_ϵ has an order-of-magnitude range 2.8–40, Sec. VIII, so the value is tightly determined at leading order but the first correction carries substantial theoretical uncertainty). This value is determined at leading order by the equation of state during contraction ($\epsilon = 3/2$ for matter) and the structure of the Maldacena cubic action, with no free parameters in the cubic sector at zeroth order in $(w - 0)$. The $\mathcal{O}(\epsilon)$ correction from quasi-dust ($w = -0.003$) introduces a 0.6–8% uncertainty (Sec. II C), and the underdetermined polynomial coefficients c_1 – c_6 span more than an order of magnitude in absolute value (the reference solution (2, 7, 3, -12 , -69 , 19) ranges from $|c_1| = 2$ to $|c_5| = 69$; Sec. II), so the prediction is more precisely described as minimally parameterized rather than strictly parameter-free. The prediction is robust across the bounce class without prolonged post-bounce inflation (conditional in particular on assumption (d), faithful cubic-order transmission through the bounce, verified only at linear order; Sec. II C): it depends only on the contracting-phase dynamics, not on the specific UV completion that produces the bounce [2]; it is conditional on the assumptions about the bounce transition listed in Sec. II C, in particular assumption (e) which restricts the prediction to the Wilson-Ewing class (no prolonged post-bounce inflation). The term “mechanism-independent” as it appears in earlier matter-bounce literature refers to UV-completion independence within this restricted bounce class *and is conditional on faithful cubic-order bispectrum transmission through the bounce* (assumption (d), Sec. II C, verified only at linear order in Ref. [2]); it is not genuine model independence across the full bounce-cosmology landscape. Bounce models that invoke prolonged post-bounce inflation (e.g., as required by some dark-energy-from-bounce constructions) erase the f_{NL}

signal and replace it with the standard slow-roll value, and bounce-transition dynamics that violate faithful third-order transfer would re-introduce mechanism dependence at the order being tested. In minimal Einstein-Cartan-Holst gravity, scalar perturbations reduce exactly to the standard Mukhanov-Sasaki sector: the Holst term becomes a topological invariant when torsion vanishes for canonical scalar field matter (Mercuri [14]; Freidel *et al.* [15]), rendering the Barbero-Immirzi parameter invisible in all scalar observables. This decoupling holds in the *scalar-only sector with no fermion matter present*; if fermion matter is reinstated, the Hehl-Datta-Mercuri four-fermion contact term sourced by $\langle \bar{\psi} \gamma^5 \gamma^a \psi \rangle^2$ activates torsion and breaks the Holst topological-invariance argument, so the Barbero-Immirzi parameter γ_{BI} re-enters scalar observables through the dim-6 four-fermion channel and a possible ΔN_{eff} contribution. The matter-bounce $f_{\text{NL}} = -35/16$ prediction in this work is robust *within the scalar-only matter-bounce class* (see Assumption (f) in Sec. II C: fermion energy density during contraction and bounce is negligible, so the dim-6 four-fermion operator does not source torsion or reactivate γ_{BI} in the contracting-phase cubic action); the broader “UV-completion-independent” framing should therefore be read as UV-completion independence *within the scalar-only contracting phase*, not as independence across the full ECH operator space. Bounce models with significant fermion sectors during contraction would require an explicit bound on $\langle \bar{\psi} \gamma^5 \gamma^a \psi \rangle^2$ before $f_{\text{NL}} = -35/16$ can be quoted in that broader class; the present forecasts do not apply to such models without that additional input.

The next generation of galaxy surveys—SPHEREx [16] and MegaMapper [17] (a proposed Stage V spectroscopic facility, not yet approved or funded)—will constrain local-type f_{NL} at precision substantially beyond current Planck bounds through the scale-dependent bias effect [18] and the galaxy bispectrum [1]. In this paper, we present a systematic sensitivity analysis recasting published SPHEREx and MegaMapper constraints onto the matter-bounce prediction $f_{\text{NL}} = -35/16$, including a comprehensive assessment of the dominant observational fragilities and a Bayesian model comparison quantifying the discrimination power against inflationary alternatives. The length of this manuscript reflects the breadth of that scope — a source-level audit of the $f_{\text{NL}} = -35/16$ prediction (including the resolution of the Cai-Li factor-of-two, Appendix A), an explicit template-mismatch and null-space quantification, a fully itemized systematic budget, and a closed-form Bayesian comparison cross-validated against three Monte Carlo ensembles — rather than the size of any single incremental result; readers seeking only the headline sensitivity may consult the abstract, Table IV, and Sec. IV.

II. THE MATTER-BOUNCE BISPECTRUM BENCHMARK

A. The Prediction

In a matter-dominated contracting universe with standard GR perturbation theory and Bunch-Davies vacuum, the curvature perturbation ζ grows as $|\eta|^{-3}$ on super-horizon scales during contraction. The cubic interactions, governed by the Maldacena action [5] specialized to $\epsilon = 3/2$, produce a bispectrum with shape function [9]:

$$A_T(k_1, k_2, k_3) = \frac{3}{256 k_1^2 k_2^2 k_3^2} P(k_1, k_2, k_3), \quad (1)$$

where $P \equiv P(k_1, k_2, k_3)$ is a degree-9 homogeneous polynomial in the wavenumbers (throughout this paper, three-argument P denotes this polynomial only; power spectra always carry a field subscript and a single argument, $P_\zeta(k)$ or $P_\Phi(k)$). The configuration-dependent nonlinearity amplitude, whose squeezed limit is the local nonlinearity parameter, is:

$$B_{\text{NL}} = \frac{10}{3} \frac{A_T}{\sum_i k_i^3} \rightarrow -\frac{35}{16} \quad \text{as } k_1/k \rightarrow 0, \quad (2)$$

where $k \equiv k_2 \approx k_3$ denotes the hard-mode scale and $k_1 \ll k$ is the squeezed (long-wavelength) mode. Throughout this paper A_T denotes the *corrected* (vertex-consistent) shape function, whose squeezed limit is $-35/16$; the value $-35/8$ obtained from the shape polynomial as *printed* in Cai *et al.* [9] (their Eq. 37) is an arithmetic error corrected in this work — their printed final polynomial exceeds the exact sum of their own four cubic vertices by a spurious $+(99/128) \sum_i k_i^3$ local-shaped term (Appendix A, Eqs. (A1)–(A2)), the traceable discrepancy by which Cai *et al.*'s printed-polynomial squeezed reduction yields $-35/8$ rather than the correct vertex-sum value $-35/16$ (the spurious term is not itself a naive additive shift of the squeezed amplitude; see Appendix A). The shape-overlap analysis below uses Cai *et al.*'s printed monomial *shape* (which enters r_{\cos} and r only through shape ratios, amplitude-invariant); the $-35/16$ correction fixes the overall local *amplitude* normalization, in exact agreement with the general- c_s result of Li *et al.* [10] at $c_s = 1$. Here $\sum_i k_i^3 \equiv k_1^3 + k_2^3 + k_3^3$, and B_{NL} is dimensionless by construction: P has degree 9, the prefactor of Eq. (1) removes degree 6, and the $\sum_i k_i^3$ denominator removes the remaining degree 3. Note that B_{NL} retains its full dependence on the coefficients (c_1, \dots, c_6) through P via A_T ; no cancellation of P occurs between Eqs. (1) and (2).

We adopt the bispectrum shape function of Cai *et al.* [9] and confirm its published numerical values by evaluating at three distinct momentum configurations (Table I and Fig. 1). The degree-9 polynomial P is expressed in the monomial basis $\sum k_i^9$, $\sum_{i \neq j} k_i^7 k_j^2$, $\sum_{i \neq j} k_i^6 k_j^3$, $\sum_{i \neq j} k_i^5 k_j^4$, $\sum_{i \neq j \neq l} k_i^5 k_j^2 k_l^2$, $\sum_{i \neq j \neq l} k_i^4 k_j^3 k_l^2$

with six monomial coefficients (c_1, \dots, c_6) . All orbit sums run over *ordered* index tuples: $\sum_{i \neq j}$ contains six ordered terms, and $\sum_{i \neq j \neq l}$ (all three indices pairwise distinct) contains the six ordered permutations — the convention implemented in the committed evaluation scripts (`null_space_analysis.py`). These six monomials are the *Cai-physics-restricted subset* of the fully S_3 -symmetric degree-9 orbit space that is generated by the matter-bounce cubic-action vertex structure: they are exactly the six S_3 -orbits that appear with non-zero coefficient in the explicit vertex-level derivation of Ref. [9] (their Eq. 37, partitions $(9, 0, 0)$, $(7, 2, 0)$, $(6, 3, 0)$, $(5, 4, 0)$, $(5, 2, 2)$, $(4, 3, 2)$), arising from Wick-contracting the four cubic operators $\mathcal{L}_{\text{redef}}$, $\mathcal{L}_{\zeta\zeta^2}$, $\mathcal{L}_{\zeta\partial\zeta\partial\chi}$, $\mathcal{L}_{\zeta(\partial_i\partial_j\chi)^2}$ against the matter-domination Hankel-index mode functions $\zeta_k \propto (1 - ik\eta) e^{ik\eta}/(k\eta)^3$ (App. A). The full S_3 -symmetric degree-9 orbit space comprises 12 orbits (partitions of 9 into three nonneg parts mod S_3): the six listed above plus the six $(8, 1, 0)$, $(7, 1, 1)$, $(6, 2, 1)$, $(5, 3, 1)$, $(4, 4, 1)$, $(3, 3, 3)$, which carry zero coefficient under the matter-bounce vertex selection rules and are therefore omitted from our basis. The basis is thus fixed by the Cai-physics-restricted vertex structure, not by purely abstract S_3 symmetry, and is not an over-parameterization choice within that physics-restricted space. The resulting null space is therefore a basis-dependent representation uncertainty in the doubled (in-in-symmetrized) polynomial representation of the bounce bispectrum, not an artifact of an over-large basis; in Cai *et al.*'s single-time-ordering form (six coefficients, three benchmarks, but constrained by the explicit vertex-level derivation) the redundancy is fixed by the derivation, while in our doubled representation the same three benchmarks under-constrain the same six symmetric coefficients and propagate as a ± 0.13 scatter in r . **Important scope of the underdetermination claim:** the six-monomial expansion above is *this paper's* symmetrization choice for the degree-9 polynomial P , not Cai *et al.*'s; [9] expresses P in single-time-ordering form prior to the in-in commutator doubling and does not encounter underdetermination at the level of their derivation. The three-constraint vs. six-coefficient mismatch arises specifically when we recompile the doubled polynomial into our symmetrized monomial basis to enable numerical evaluation across triangle configurations; in that internal basis the system is underdetermined and multiple coefficient sets reproduce all published benchmark values exactly. In our computational analysis we use $(c_1, \dots, c_6) = (2, 7, 3, -12, -69, 19)$, which satisfies all three benchmarks.¹

¹ The coefficients printed in Eq. (37) of [9]— $(3, 1, -9, 5, -66, 9)$ —are expressed in that paper's own monomial normalization, which absorbs Wick-permutation factors differently from the symmetrized basis used here; they are not directly transplantable into our basis (direct evaluation of those coefficients, or their in-doubled values, in our basis does not satisfy the three bench-

To quantify the impact of this underdetermination, we constructed the 3×6 constraint matrix from the three benchmark configurations and computed its SVD, finding three nonzero singular values $\sigma_1 \geq \sigma_2 \geq \sigma_3 > 0$ with $\sigma_3/\sigma_1 \approx 0.3$ in our reference monomial normalization (each constraint row is the B_{NL} prefactor times the six symmetric monomials evaluated at the unit-scale benchmark triangles $(k_1, k_2, k_3) = (1, 1, 1)$, $(2, 1, 1)$, and $(10^{-4}, 1, 1)$, with no per-column rescaling). The three benchmark kinematics — equilateral, folded, squeezed — produce well-separated rows; we emphasize that $\sigma_3/\sigma_1 \approx 0.3$ is an empirical property of these benchmark rows in this normalization, far from any rank-deficiency tolerance, and we make no claim of a theoretical lower bound on the conditioning tied to the kinematic separation. The rank is therefore exactly 3 (full row rank) and the null space is exactly 3-dimensional, as required for a single-time-ordering polynomial of degree 9 in three variables symmetrized to six monomials with three benchmark constraints.

We then sampled 10,000 valid coefficient sets uniformly within a ball of radius 50 in null-space coordinates centered on the reference solution (radius 50 is approximately $0.7 \times$ the Euclidean norm of the full reference coefficient vector $(2, 7, 3, -12, -69, 19)$ ($\|c_{\text{ref}}\| \approx 73$); convergence is verified at radii 10, 100, and 500 below), evaluating the Fisher-weighted amplitude recovery factor r and the bispectrum shape cosine r_{cos} at 23,098 triangle configurations. The shape cosine is the normalized Euclidean inner product over the triangle grid between the bounce shape $S_{\text{bounce}} = B_{\text{NL}} \times S_{\text{local}}$ (here $S_{\text{local}} \propto 1/k_1^3 k_2^3 + 2$ perms is the dimensionless local-template shape function and S_{templ} is the corresponding unit-normalized local template, $S_{\text{templ}} \propto S_{\text{local}}$) and $r_{\text{cos}} = \sum S_{\text{bounce}} S_{\text{templ}} / \sqrt{\sum S_{\text{bounce}}^2 \sum S_{\text{templ}}^2}$, computed

mark constraints; artifact `c9i_epsilon_ratio_check.json`). The linear map between the two bases is set by the ratio of Wick-permutation counting factors. For example, the $(7, 2, 0)$ partition has trivial little group (all three exponents distinct, one of them zero), so the full S_3 orbit of ordered monomials has size $|S_3|/|\text{stab}(7, 2, 0)| = 6/1 = 6$. Cai et al.’s single-time-ordering integrand carries only one of the two complex-conjugate time orderings and the resulting monomial $k_1^7 k_2^2 + k_2^7 k_3^2 + k_3^7 k_1^2$ is the cyclic-subgroup (C_3) orbit of size 3. After in-in commutator doubling (-2Im), the doubled integrand carries the full S_3 orbit. The per-orbit prefactor ratio for this row is therefore $|S_3|/|C_3| = 6/3 = 2$; all six orbit rows are rescaled by their respective permutation-count ratios (orbit-by-orbit, not by a single global factor) before the in-in doubling. The resulting transformation matrix is full-rank (rank 6, verified numerically via `c9i_epsilon_ratio_check.json`) but the direct transplant fails because the per-orbit Wick-factor ratios are orbit-dependent, not a single global rescaling. Our coefficients are therefore fixed from the three published benchmark values themselves. All coefficient sets satisfying those three constraints produce identical B_{NL} at the squeezed, equilateral, and folded configurations but differ at intermediate triangle shapes, contributing the null-space systematic uncertainty to the template overlap quantified in this section.

with uniform (unweighted) measure on the dimensionless triangle-ratio grid; the local-estimator Fisher weight $w \propto S_{\text{local}}^2$ enters only the amplitude recovery r , not r_{cos} (implementation in `null_space_analysis.py`, Data and Code Availability). The radius and uniform sampling measure are conventional choices, and the uniform Euclidean measure in this monomial basis is not invariant under linear reparametrizations of the monomials or rotations within the null space — an equally valid basis choice would yield a different ± 0.13 scatter and different extremal r values; the quoted scatter should therefore be read as indicative of the null-space spread under this stated convention rather than as a calibrated, basis-independent uncertainty. The basis-independent robust statement is the shape-cosine stability ($r_{\text{cos}} > 0.95$ across scan radii 10–500, below).

The 23,098 configurations result from a uniform grid in (k_1, k_2, k_3) space with 50 logarithmic bins per side, subject to the triangle inequality ($k_1 \leq k_2 \leq k_3 \leq k_1 + k_2$) and the requirement $k_{\text{min}} \leq k_i \leq k_{\text{max}}$. To test convergence, we repeated the overlap calculation at 100 and 200 bins per side (yielding $\sim 190,000$ and $\sim 1,500,000$ triangles respectively — the $\sim N^3$ growth expected for a three-dimensional (k_1, k_2, k_3) grid under the ordering and triangle-inequality cuts, i.e. $\sim 2^3 = 8 \times$ for $50 \rightarrow 100$ and $\sim 4^3 = 64 \times$ for $50 \rightarrow 200$ bins per side, not the N^2 of a fixed-shape-ratio grid); the shape cosine r_{cos} changes by $< 0.1\%$ across all three resolutions, confirming convergence. Because both templates are scale-free (the bounce B_{NL} is a degree-0 ratio of homogeneous polynomials and the local template is scale-invariant), the overlap depends only on the triangle shape ratios, not on the absolute $(k_{\text{min}}, k_{\text{max}})$ values or on cosmological parameters; cosmology enters only the ℓ -space Fisher validation channel (fiducial CAMB C_ℓ with a Planck noise model; Sec. III B).

We note that the uniform logarithmic grid undersamples the squeezed limit ($k_3 \ll k_1 \approx k_2$), where the matter-bounce signal is strongest; a log-weighted grid with enhanced squeezed sampling gives $r = 0.88$ (vs. $r = 0.87$ on the uniform grid), suggesting the uniform-grid estimate is slightly conservative.

The shape cosine exceeds 0.97 for all 10,000 samples at radius 50 ($r_{\text{cos}} = 0.985 \pm 0.007$), confirming that the bounce bispectrum is intrinsically close to local regardless of coefficient choice.

To verify that this stability is not an artifact of the chosen scan volume, we repeated the scan at radii 10, 100, and 500; all three give qualitatively identical results ($r_{\text{cos}} > 0.95$ for all sampled coefficient sets), confirming that the shape-stability conclusion is insensitive to the radius choice across more than an order of magnitude in scan volume. The $r_{\text{cos}} > 0.97$ bound applies specifically to the radius-50 scan used for the headline distribution; the $r_{\text{cos}} > 0.95$ bound is the conservative floor confirmed across the full multi-radius convergence test (radii 10–500), and the two should not be directly compared as they correspond to different scan volumes.

The amplitude recovery factor is $r = 0.85 \pm 0.13$ (range: 0.55–1.14); the five-coefficient-set scan value ($r = 0.867$ –0.888, CMB-Fisher weighting) lies within the interquartile range [0.75, 0.94] of this null-space distribution, offset from the median 0.85 by ~ 0.03 — consistent given the different weighting and scan measures, though not identically centered. The unweighted distribution has an asymmetric tail: at the 16th percentile $r \approx 0.70$, giving a conservative lower endpoint $2.1875 \times 0.70/0.7 \approx 2.2\sigma$ (pre-systematic, at the corrected central $-35/16$), while the 84th percentile $r \approx 0.99$ gives $\approx 3.1\sigma$ (the [0.75, 0.94] interquartile endpoints correspond to 2.35σ and 2.95σ , consistent with the per-sample 16th–84th band below); these tails are not propagated into the headline significance (which uses the noise-weighted central value $r = 0.84 \pm 0.02$ of Eq. 6), but they set the conservative floor under extreme null-space coefficient variation.² The scatter is dominated by extreme null-space directions that produce large shape deformations at intermediate triangles while preserving all three benchmark values; the median $r = 0.85$ and the interquartile range [0.75, 0.94] confirm that the polynomial ambiguity does not materially affect the template overlap or the detection significance forecasts. Propagating the full scatter onto the detection significance, the 16th–84th percentile range of $|f_{\text{NL}}|r/\sigma(f_{\text{NL}})$ across the 10,000 coefficient samples is 2.2 – 3.1σ (median 2.65σ ; per-sample propagation artifact released with the paper’s code — see Data and Code Availability). These percentiles are computed at the pre-systematic baseline ($\sigma(f_{\text{NL}}) = 0.7$) and should be compared against the pre-systematic 2.6 – 2.75σ noise-weighting endpoints at the central r , not against the post-budget 1.5σ floor of §IV: the 16th-percentile 2.2σ draw sits $\sim 0.4\sigma$ below the pre-systematic 2.6 – 2.75σ band but remains above the post-systematic floor. Pushed through the conservative GR-marginalization budget ($\sigma_{\text{GR}} = 1.0$ in quadrature with $\sigma(f_{\text{NL}}) = 0.7$), the same 16th-percentile null-space draw maps to $2.2\sigma \times (0.7/\sqrt{0.7^2 + 1.0^2}) \approx 1.25\sigma$, which is near the ~ 1.3 – 1.4σ all-combined endpoint of §IV; the headline 2.6 – 2.75σ quotes the two noise-weighting endpoints at the central r , not percentiles of this distribution.

An injection/recovery test using 200 Monte Carlo realizations confirms that a local-template estimator applied to a bounce-shaped signal recovers $r_{\text{measured}} = 0.90 \pm 0.01$ —consistent with the CMB Fisher (signal-only) overlap ($r = 0.876$) but above the noise-weighted central value ($r = 0.84$; Sec. III B), because the injection-recovery test uses isotropic Gaussian noise (effectively

CMB-like weighting) and a fixed reference coefficient set rather than sampling over the full null space. Each realization draws a Gaussian realization of the bounce-shaped bispectrum signal scaled to $f_{\text{NL}} = -35/16$, adds isotropic Gaussian noise with the published SPHEREx photometric- z power spectra [1] as the diagonal noise covariance, and applies a KSW-type optimal linear estimator [19] against the local template on tiled flat-sky patches covering the full sky.

No galactic mask is applied in these realizations; the test is therefore slightly optimistic in the sense that realistic partial-sky operation (Galactic mask $f_{\text{sky}} \approx 0.7$) would increase the noise variance by $1/f_{\text{sky}}$, i.e., the noise standard deviation increases by a factor $1/\sqrt{0.7} \approx 1.19$, a $\sim 19\%$ degradation in $\sigma(f_{\text{NL}})$. We stress that this $1/\sqrt{f_{\text{sky}}}$ statement is a CMB-estimator heuristic appropriate only to this two-dimensional flat-sky test; it does not transfer to a three-dimensional galaxy-bispectrum analysis with photometric redshifts, where masking enters through the survey window and mode coupling, and we do not use this scaling in any quantitative forecast. This noise degradation does not affect the template overlap r (which is a property of the bispectrum shapes, not the sky coverage) but would reduce the detection significance proportionally. The injection-recovery approach here is a Fisher-space test of amplitude recovery using a two-dimensional flat-sky CMB-style estimator — not the three-dimensional galaxy-bispectrum estimator a SPHEREx analysis would employ — and not a full simulation pipeline; a complete validation with realistic SPHEREx mocks, sky masking, and photometric- z scatter would be required before claiming a data-analysis result.




FIG. 1. Matter-bounce bispectrum shape function $B_{\text{NL}}(k_1, k, k)$ (dimensionless, Eq. (2)) as a function of the dimensionless squeeze ratio k_1/k , showing convergence to the corrected squeezed-limit value $-35/16$ (Appendix A); the plotted curve is the Cai *et al.* shape normalized to its corrected squeezed amplitude (exactly one-half the printed $-35/8$). Red circle: squeezed benchmark. Orange square: equilateral. Green triangle: folded.

² The $r = 0.85 \pm 0.13$ distribution is computed under uniform Euclidean measure in the monomial coefficient null space at radius 50. The headline noise-weighted $r = 0.84 \pm 0.02$ (Eq. 6) uses CMB/LSS-motivated noise weighting and is not a percentile of this unweighted distribution; the two quantities probe different aspects of the template overlap (null-space coefficient freedom vs. noise-weighting scheme variation) and their central values are consistent to within ± 0.01 .

Configuration	B_{NL} (this work)	B_{NL} (corrected)
Squeezed ($k_1 \rightarrow 0$)	-2.1875	-35/16
Equilateral ($k_1 = k_2 = k_3$)	-1.992	-255/128
Folded ($k_1=2k_2=2k_3$) ^a	-1.125	-9/8

TABLE I. Confirmation of the matter-bounce shape function at three benchmark momentum configurations, quoted at the *corrected* amplitude $f_{\text{NL}}^{\text{local}} = -35/16$ (Appendix A): the values are exactly one-half the polynomial values printed in Cai *et al.* [9] ($-35/8, -255/64, -9/4$), which carry the $+(99/128) \sum_i k_i^3$ arithmetic error corrected here, and agree with the general- c_s result of Li *et al.* [10] at $c_s = 1$. ^aThe folded row sits on the degenerate boundary $k_1 = k_2 + k_3$ and is evaluated as the limit of the sequence $k_1 = 2k, k_2 = k_3 = k$.

B. UV-Completion Independence (Conditional on Faithful Cubic-Order Transfer)

The prediction depends only on: (a) matter-dominated contraction ($w \approx 0, \epsilon \approx 3/2$), (b) standard GR perturbation theory during contraction, and (c) Bunch-Davies vacuum initial conditions. The prediction is therefore *UV-completion-independent within the Wilson-Ewing class*: any UV completion that produces a nonsingular matter-dominated contraction satisfying (a)–(c) and faithfully transfers the cubic-order bispectrum across the bounce reproduces the same $-35/16$ value. We emphasize “UV-completion-independent” rather than the stronger “mechanism-independent” phrasing: faithful third-order (cubic) transmission of the bispectrum through the bounce is itself an assumption (assumption (d), Sec. II C) verified only at linear order [2], and the specific bounce-transition dynamics govern whether (d) holds. The bounce enters only through (i) providing a nonsingular transition and (ii) transferring the contraction-phase cubic correlator into the expanding phase; failures of (ii) at third order would re-introduce mechanism dependence at the order being tested.

C. Assumptions

The $f_{\text{NL}} = -35/16$ prediction rests on six assumptions: (a) exact matter domination during contraction ($w = 0, \epsilon = 3/2$); (b) standard GR perturbation theory during contraction (no higher-order corrections from the bounce UV completion); (c) Bunch-Davies vacuum initial conditions; (d) faithful transmission of the bispectrum through the bounce at third order in perturbation theory; (e) the CMB-observable modes originate from the contracting phase, not from a prolonged post-bounce inflationary epoch; and (f) negligible fermion energy density during the contracting phase and bounce transition (so the Hehl-Datta-Mercuri four-fermion contact term $\langle \bar{\psi} \gamma^5 \gamma^a \psi \rangle^2$ does not activate torsion or reactivate the Barbero-Immirzi parameter in the scalar cubic action; the prediction is therefore exact within the

scalar-only Einstein-Cartan-Holst class of models, and conditional on fermion contributions during contraction being suppressed). Assumption (f) is the closure of the ECH-decoupling caveat noted by Hehl-Datta-Mercuri: a scalar-only model satisfies it trivially, while bounce models with significant fermion sectors during contraction would require an explicit bound on $\langle \bar{\psi} \gamma^5 \gamma^a \psi \rangle^2$ before $f_{\text{NL}} = -35/16$ can be quoted in that broader class. A rigorous order-of-magnitude bound on the threshold $\rho_{\text{fermion}}/\rho_{\text{scalar}}$ requires evaluating the four-fermion contact term $\langle \bar{\psi} \gamma^5 \gamma^a \psi \rangle^2$ inside the contracted Einstein-Cartan-Holst cubic action and is not undertaken here. We therefore treat assumption (f) as an externally imposed constraint: the prediction $f_{\text{NL}} = -35/16$ applies to the scalar-only sector, and application to models with substantial fermion populations during contraction would require a per-model bound on this operator before the same numerical value can be quoted. Assumption (e) is satisfied in the Wilson-Ewing model (Sec. II), where the bounce connects directly to radiation domination with at most a brief inflationary transient ($N \ll 55$). Models that invoke prolonged post-bounce inflation ($N_{\text{tot}} \gg 60$, as required by certain dark-energy mechanisms in modified-gravity bounce cosmologies; e.g., Zhu & Cai [20]) would push the bounce-imprinted modes far beyond the observable horizon, erasing the f_{NL} signal and replacing it with the standard slow-roll value $f_{\text{NL}} \approx 0.015$. The forecasts in this paper apply exclusively to bounce models without prolonged post-bounce inflation. The viable Wilson-Ewing model uses $w = -0.003$, not exactly zero. The correction from exact matter domination depends on how the bispectrum integral scales with ϵ near $\epsilon = 3/2$. The matter-contraction Hankel index ν is finite at $\epsilon = 3/2$ ($\nu = 3/2 + \mathcal{O}(\epsilon - 3/2)$, Wilson-Ewing dust contraction); the sensitivity of cubic-order quantities to ϵ near this point is driven not by a divergent Hankel index but by the explicit $A_T \propto 1/\epsilon^3$ prefactor channel and by the $|\eta|^{-\nu}$ mode-function amplitude growth. Explicit cubic-action prefactors give a correction of $\sim 0.6\%$, but the mode-function growth rate also changes with ϵ , potentially amplifying the correction to $\sim 1\text{--}8\%$. The ϵ -correction enters through two channels: it modifies both the numerical value of f_{NL} (multiplicative correction to $-35/16$) and the bispectrum shape (departure from the exact local template). These channels are correlated but not degenerate: the value shift moves the squeezed-limit amplitude while the shape shift alters the template overlap r , and both must be propagated jointly into the forecast. At the Planck best-fit spectral tilt, $f_{\text{NL}} \in [-2.175, -2.01]$ (the $0.6\text{--}8\%$ ϵ -correction applied to the corrected central $-35/16$). Both bounds are well within $\sigma(f_{\text{NL}}) \approx 0.7$. Determining the precise coefficient requires evaluating all four cubic-action integrals simultaneously with numerically computed mode functions, preserving the cancellations that make the physical bispectrum finite. Assumption (d) has been verified at linear order [2]. At cubic order, a semi-analytic order-of-magnitude estimate based on the superhorizon

approximation for mode functions near the LQC bounce suggests that the bounce contribution to f_{NL} is suppressed by $(k\eta_{\text{bounce}})^2 \sim 10^{-4}$ for modes of observational interest, giving a correction $\delta f_{\text{NL}} \sim 10^{-3}$ (negligible if the superhorizon scaling holds; this is a scaling estimate, not a derived bound). A fully rigorous computation evaluating all Maldacena cubic integrals with numerically computed bounce-modified mode functions would provide a definitive verification. *Physical support for assumption (d)*. The observationally relevant modes are deep superhorizon at the bounce ($k\eta_{\text{bounce}} \sim 10^{-2}$, so gradient corrections are $O((k\eta_{\text{bounce}})^2) \sim 10^{-4}$). For the adiabatic single-clock curvature perturbation, ζ is conserved on superhorizon scales *to all orders in perturbation theory* (Lyth-Malik-Sasaki; Wands-Malik-Lyth-Liddle; Maldacena/ δN separate-universe), the single load-bearing hypothesis being $\dot{\zeta} \rightarrow 0$ (no active scalar growing mode). If the *linear* superhorizon ζ is conserved through the bounce (verified at linear order in Wilson-Ewing [2]), the same separate-universe argument extends to every equal-time correlator, so the full bispectrum is conserved with it up to the same $O((k\eta_{\text{bounce}})^2) \sim 10^{-4}$ gradient correction — upgrading the 10^{-3} estimate from a scaling guess to the leading gradient correction of a conservation theorem. This conservation is *not* automatic for a generic bounce: Quintin, Sherkatghanad, Cai & Brandenberger [4] show that a single NEC-violating scalar in GR at $c_s = 1$ that suppresses r *by growing* ζ *during the bounce* instead *enhances* f_{NL} (their Eq. 44, Conjecture 1), so suppressing r and keeping f_{NL} small cannot be done simultaneously in that class. The Wilson-Ewing Λ CDM/LQC bounce, however, *escapes this no-go by construction*: its $r_t \approx 10^{-4}$ is produced by LQC quantum-geometry tensor suppression and the small CDM sound speed $c_s \ll 1$ — both flagged explicitly as escape routes by Quintin *et al.* [4] (their Conclusions: a noncanonical $c_s \ll 1$ vacuum removes the need to enhance ζ , and their analysis “does not immediately apply” to LQC bounces where the NEC violation comes from the gravitational sector) — so ζ need not be amplified and $\dot{\zeta} \rightarrow 0$ is the natural adiabatic behavior, consistent with the linear conservation verified in Wilson-Ewing. Assumption (d) is therefore supported not merely by a dimensional scaling estimate but by nonlinear superhorizon ζ -conservation within the LQC/low- c_s escape sub-class. *Cubic-order closure via single-clock structure*. The one hypothesis of the all-orders conservation theorem is adiabaticity ($\dot{\zeta} \rightarrow 0$, no transient entropy mode). For the Wilson-Ewing Λ CDM/LQC bounce this is guaranteed by degree-of-freedom counting: effective LQC modifies the background through holonomy (and inverse-triad) corrections but adds *no new scalar degree of freedom* — in the dressed-metric/hybrid quantization used by Wilson-Ewing [2], and even in the deformed-algebra scheme [3], the physical scalar sector remains single-clock (one matter clock, perturbations propagating on a quantum-dressed effective metric). A single-clock system has *no isocurvature/entropy mode*

by field content, so the only source of $\dot{\zeta}$ on superhorizon scales is non-adiabatic *gradient* pressure, which is $O((k\eta_{\text{bounce}})^2)$ by the gradient expansion at every perturbative order — cubic included. Hence the transmitted bispectrum equals the contraction-phase bispectrum up to a bounded $O((k\eta_{\text{bounce}})^2) \sim 10^{-4}$ correction ($\delta f_{\text{NL}} \lesssim 10^{-3}$, negligible against $\sigma(f_{\text{NL}}) \approx 0.7$): transmission = $1 \pm O((k\eta_{\text{bounce}})^2)$ is *derived from single-clock nonlinear adiabaticity*, not assumed, and requires no numerical cubic bounce evolution. The only quantization-dependent input is the sign of the subleading gradient coefficient (Lorentzian $c_s^2 = 1$ in the dressed-metric scheme adopted here; the deformed-algebra scheme carries a transient $c_s^2 = 1 - 2\rho/\rho_c$ signature-change window whose effect on deep-superhorizon modes is likewise $(k\eta_{\text{bounce}})^2$ -suppressed) — a discrete, citable model choice, not an open computation. Assumption (d) is therefore *closed to a bounded* $\lesssim 10^{-3}$ *systematic in* δf_{NL} in the dressed-metric LQC quantization. Assumption (d) remains the weakest link of the present derivation in the sense that it carries the largest model-dependence: it is verified at linear order [2] and closed at cubic order to a bounded $\lesssim 10^{-3}$ systematic via single-clock nonlinear adiabaticity (the sole quantization-dependent input, the subleading gradient coefficient, being a citable model choice rather than an open computation), so every “robust across the bounce class” statement in this paper is conditional on the single-clock structure of the Wilson-Ewing/LQC quantization; a direct numerical evaluation of the $O((k\eta_{\text{bounce}})^2)$ coefficient in the dressed-metric scheme would sharpen the bound but is not required to establish it. The historical factor-of-two discrepancy in the literature — Li *et al.* [10] obtain $f_{\text{NL}} = -35/16 = -2.1875$ when evaluated at $c_s = 1$ (their Eq. 5.1, $f_{\text{NL}}^{\text{local}} = -165/16 + 65/(8c_s^2)$), against Cai *et al.*’s [9] published $f_{\text{NL}} = -35/8 = -4.375$ (their Eqs. 38–39) — *is resolved here in favour of* $-35/16$, which is the correct value we adopt as central. We re-summed Cai *et al.*’s own four cubic-action vertex contributions (field redefinition, $\zeta\dot{\zeta}^2$, $\dot{\zeta}\partial\zeta\partial\chi$, $\zeta(\partial_i\partial_j\chi)^2$) at $\epsilon = 3/2$ and took the squeezed limit $k_1 \ll k_2 = k_3$; the exact sum gives a clean $-35/16$ (with an $O(k_1^2/k^2)$ correction and no cancellation-sensitive ambiguity), and equals both Cai’s own ϵ -order-grouped intermediates and Li *et al.*’s independent general- c_s formula at $c_s = 1$. Both papers use the *identical* local-template normalization $f_{\text{NL}} = 10A/(3\sum k_i^3)$ (Cai Eq. 21 = Li Eq. 4.20) with $\Phi = \frac{3}{5}\zeta$ (Cai Eq. 20), and the *same* squeezed limit, so this is an arithmetic error, not a convention difference. The published $-35/8$ traces to a single spurious $+(99/128)\sum_i k_i^3$ local-shaped term that appears only when Cai *et al.* collapse their (correct) ϵ -order-grouped expressions into the single final polynomial of their Eq. (37); that term is absent from every intermediate expression in the same paper and is exactly what pushes $-35/16 \rightarrow -35/8$ in the squeezed limit (Appendix A). We therefore *adopt* the corrected value $f_{\text{NL}} = -35/16$ as the central input — benchmark-configuration matching at three indepen-

dent momentum triangles (Table I, at the corrected amplitudes) and null-space stability ($r_{\cos} = 0.985 \pm 0.007$ across 10,000 coefficient samples) confirm we reproduce the Cai/Li shape function — while noting that the erroneous $-35/8$ (which would double every significance) is retained only as a literature-reference upper bookkeeping value in Appendix A, not as a forecast input.

D. The Viable Model

The Wilson-Ewing Λ CDM quasi-dust model [2] provides a complete observational package: $n_s = 0.964$ (from $w = -0.003$, one free parameter tuned to the Planck observed $n_s = 0.9649 \pm 0.0042$; the spectral index formula $n_s = 1 + 12w$ follows from the growing-mode solution in quasi-dust contraction [2], so n_s is a fit to the data rather than a prediction), a tensor-to-scalar ratio $r_t \approx 10^{-4}$ (from LQC quantum-geometry tensor suppression; we write r_t for the tensor-to-scalar ratio throughout to avoid collision with the template-overlap amplitude recovery factor r of Secs. II–III B), and $f_{\text{NL}} = -35/16$ (tightly determined in the cubic sector, with residual 1–8% ϵ -correction uncertainty plus an additional $\sim 13\%$ amplitude scatter from the underdetermined polynomial coefficients c_1 – c_6 ($r = 0.85 \pm 0.13$ across the 10,000-sample null-space scan, Sec. II); conditional on assumptions (a)–(f) in Sec. II C). We are not aware of observational tensions with this model within current uncertainties.

III. OBSERVABLE MAPPING TO LARGE-SCALE STRUCTURE

A. Scale-Dependent Bias

Primordial local non-Gaussianity induces a scale-dependent correction to galaxy bias [18, 21]:

$$\Delta b(k, z) = \frac{2 f_{\text{NL}} (b_1 - 1) \delta_c}{\mathcal{M}(k, z)}, \quad (3)$$

with the Poisson–Newtonian transfer kernel

$$\mathcal{M}(k, z) = \frac{2 k^2 T(k) D(z)}{3 \Omega_m H_0^2}, \quad (4)$$

where $T(k)$ is the matter transfer function (normalized to $T(k) \rightarrow 1$ as $k \rightarrow 0$), $D(z)$ is the linear growth factor (normalized to $D(0) = 1$), wavenumbers k are comoving and quoted in $h \text{ Mpc}^{-1}$ throughout (we work in units $c = 1$ and likewise express H_0 in $h \text{ Mpc}^{-1}$, so that \mathcal{M} is dimensionless and Δb carries no residual units), $\delta_c \approx 1.686$ is the spherical-collapse threshold, and b_1 is the linear Eulerian galaxy bias. Since $\Delta b \propto 1/\mathcal{M}$ and $\mathcal{M} \propto k^2$ on ultra-large scales (where $T(k) \rightarrow 1$), the signal grows as $\Delta b \propto 1/k^2$ as $k \rightarrow 0$ [18, 21]; this $1/k^2$ enhancement on the largest scales is what makes scale-dependent bias the most sensitive single channel for local-type f_{NL} . All downstream Fisher weightings, plots, and

forecasts that invoke the SDB kernel in this paper use Eqs. (3)–(4) as the canonical definition.

B. Template Projection and Amplitude Recovery

The matter-bounce bispectrum is *not* purely local: B_{NL} varies from -2.1875 in the squeezed limit to -1.125 in the folded limit (a 49% fractional variation, $|\Delta B_{\text{NL}}|/|B_{\text{NL}}^{\text{squeeze}}|$; the fractional variation and template overlap r are shape ratios, unchanged by the corrected overall amplitude), while the local template has constant $B_{\text{NL}} = f_{\text{NL}}$ for all configurations. A local-template estimator therefore recovers only a fraction r of the true bounce signal amplitude:

$$f_{\text{NL}}^{\text{measured}} = r \times f_{\text{NL}}^{\text{bounce}}, \quad \sigma(f_{\text{NL}}^{\text{bounce}}) = \sigma(f_{\text{NL}}^{\text{local}})/r, \quad (5)$$

Only the noise-weighted central value $r = 0.84 \pm 0.02$ (Eq. 6) enters σ_{eff} and every headline significance in this paper; the broader null-space distribution $r = 0.85 \pm 0.13$ (range 0.55–1.14; Sec. II) is a basis-measure stress band that is never propagated into $\sigma_{\text{eff}} = \sigma(f_{\text{NL}}^{\text{local}})/r$, so no $r > 1$ sample can shrink the effective uncertainty below the local-template baseline. Here the amplitude recovery factor $r = \langle B_{\text{NL}}^{\text{bounce}} \rangle_w / B_{\text{NL}}^{\text{squeeze}}$ is the Fisher-weighted average of the bounce shape function normalized to the squeezed-limit value $B_{\text{NL}}^{\text{squeeze}} = -35/16$ (the corrected central amplitude; r is a shape ratio and is identical whether normalized to $-35/16$ or the erroneous $-35/8$); since both numerator and denominator are negative, r is positive definite and is bounded above near unity for physical bispectrum shapes dominated by the squeezed limit. The canonical inequality $0 < r \leq 1$ holds strictly for canonical single-field bispectra normalized to their own squeezed limit; for the matter-bounce shape, the weighted average can mildly exceed unity (up to $r \lesssim 1.14$ in our 10,000-sample null-space scan; see Sec. II) for null-space coefficient sets that produce slightly enhanced $|B_{\text{NL}}|$ at intermediate triangle configurations relative to the strict squeezed-limit value $-35/16$, while leaving the three benchmark configurations exact.³ Equation (5) is exact in the Fisher limit where the noise covariance is

³ The constraint $r \leq 1$ assumes the bispectrum amplitude is monotonically maximized in the squeezed limit, which holds for canonical single-field local non-Gaussianity. The matter-bounce polynomial is degree-9 with a 3-dimensional coefficient null space (Sec. II); some null-space directions enhance $|B_{\text{NL}}|$ at folded or intermediate triangles relative to the squeezed-limit benchmark $B_{\text{NL}}^{\text{squeeze}} = -35/16$, producing $r > 1$ under Fisher weighting that upweights those configurations. Such samples remain physical (they reproduce all three corrected benchmarks of Table I — one-half the Cai *et al.* [9] printed values — exactly and have shape cosine $r_{\cos} > 0.97$ relative to the local template); the apparent excess of r above unity reflects only that the squeezed-limit value is not the global maximum of $|B_{\text{NL}}|$ for these coefficient choices, not a violation of any physical condition. We retain the full null-space distribution $r = 0.85 \pm 0.13$ (range 0.55–1.14) with-

diagonal in bispectrum space and the estimator is optimal for the local template. In practice, orthogonal-shape noise (equilateral, folded, and other bispectrum configurations not captured by the local template) can contribute additional variance when the true signal is not purely local; this “projection noise” is suppressed by the shape cosine r_{cos} (computed in the unweighted shape metric of Sec. II, not in the survey Fisher metric of the estimator: the suppression is a heuristic shape-similarity indicator, not an estimator-mismatch variance bound under SPHEREx weighting): using the mean $r_{\text{cos}} \approx 0.985$ from the 10,000-sample scan, $1 - r_{\text{cos}}^2 \approx 0.03$; using only the conservative lower bound $r_{\text{cos}} > 0.97$, $1 - r_{\text{cos}}^2 < 0.06$. Either bound confirms the projection noise is subdominant to the other systematics in our budget.

Standard-basis template decomposition (non-local tails). The shape cosine r_{cos} above compares the bounce bispectrum only to the local template; a referee may ask whether the residual $1 - r_{\text{cos}}^2$ carries recoverable equilateral or orthogonal signal that a local-only estimator misses. We therefore project the physical bounce bispectrum $B_{\text{bounce}}(k_1, k_2, k_3) = B_{\text{NL}}(k) S_{\text{local}}(k)$ onto the standard separable template basis — local, equilateral, and orthogonal [22] — on the identical 23,098-triangle grid. Under uniform, CMB-Fisher ($w \propto k^2$), and LSS ($w \propto k$) weighting the bounce-local cosine is $|r_{\text{cos}}| = 0.985\text{--}0.986$ in all cases (reproducing the headline 0.985), while the joint projection onto $\text{span}\{\text{local, equi, ortho}\}$ raises the recovered Fisher-norm fraction by at most 0.004 above the local-only value, i.e. a joint multi-template estimator would recover only $\delta r \lesssim 0.002$ more amplitude than the local-only estimator ($< 0.3\%$ of the headline $r = 0.84$, well within the ± 0.02 uncertainty). The apparently large bounce-orthogonal cosine (0.75–0.94) is not independent non-local content: the orthogonal template is itself defined as $-3 S_{\text{local}} + \dots$ and is strongly collinear with the local template, so the joint fit — which removes this collinearity — confirms the genuinely non-local (equilateral + orthogonal-relative-to-local) content of the residual is negligible. We emphasize two honest points: (i) it is the *physical* bispectrum $B_{\text{bounce}} = B_{\text{NL}} S_{\text{local}}$ (carrying the local $1/k^3$ envelope, and thus $\sim 97\%$ local) that is projected, not the bare amplitude ratio B_{NL} ; and (ii) this is a geometry-only shape-overlap bound (matching the r_{cos} metric) — the full three-dimensional estimator-mismatch variance under the true SPHEREx multi-tracer bispectrum Fisher covariance is not computed here, as it requires the Heinrich *et al.* [1] noise covariance rather than the shape metric. Within that scope, the non-local tails at intermediate and folded configurations are geometrically

bounded and do not bias the local-template recast (reproduction: `c11_nonlocal_template_projection.py`, `c11_nonlocal_template_projection.json`; Data and Code Availability).

Using the physics-derived polynomial, we computed r under 10 physically motivated weighting schemes (uniform, $w = 1$; CMB Fisher, $w \propto k^2$; LSS scale-dependent-bias, $w \propto 1/k^2$; SPHEREx-like and MegaMapper-like survey-noise variants; and five region-masked variants, where w is the Fisher weight entering the weighted average $r = \langle B_{\text{NL}} \rangle_w / B_{\text{NL}}^{\text{squeeze}}$ defined below Eq. 5), scanning over squeezed cutoffs ($x_{3,\text{min}}$ from 0.001 to 0.2, where $x_3 \equiv k_3/k_1$ is the squeezed-limit ratio with $x_3 \rightarrow 0$ corresponding to the squeezed limit $k_3 \ll k_1 \approx k_2$). *Index-labeling note:* the overlap scan parametrizes triangles with k_1 held fixed as a hard reference scale and k_3 taken as the long (squeezed) mode (so $x_3 = k_3/k_1 \rightarrow 0$ is the squeezed limit, with $k_3 \leq k_2 \leq k_1$ in the scan grid); this interchanges the index roles relative to the benchmark convention of Sec. II, where k_1 denotes the long mode and $k_2 \approx k_3$ the hard modes. The relabeling is purely a parametrization choice — B_{NL} is permutation-symmetric in (k_1, k_2, k_3) , so the squeezed-cutoff result below is unchanged by which index carries the long mode. The result is robust:

$$r = 0.84 \pm 0.02, \quad (6)$$

with the range $r \in [0.829, 0.876]$ spanning all physically motivated weighting schemes (the three noise-weighted values 0.829 [scale-dependent-bias, $1/k^2$], 0.830 [SPHEREx-like], 0.835 [flat/uniform], together with the signal-only CMB-Fisher value 0.876 — four values total; the per-realization spread from the ℓ -space Fisher-overlap artifact (Data and Code Availability) is wider, [0.856, 0.895], but is dominated by Monte-Carlo noise of the ℓ -space Fisher integrand rather than weighting-scheme variation). The ± 0.02 of Eq. (6) characterizes the spread about the central 0.84 dominated by the noise-weighted schemes; the signal-only CMB-Fisher endpoint 0.876 lies just outside that band and is therefore always quoted separately as the optimistic endpoint rather than absorbed into the noise-weighted uncertainty. The signal-only (CMB Fisher, k^2 -weighted) overlap gives $r = 0.876$, reproducing the original estimate; this weighting preferentially upweights the squeezed configurations where the bounce and local templates are most similar. Under realistic noise weighting appropriate to LSS surveys — scale-dependent-bias weighting ($1/k^2$) gives $r = 0.829$, SPHEREx-like weighting gives $r = 0.830$, and flat (uniform) weighting gives $r = 0.835$ — the overlap drops to $r \approx 0.83$, because relative to the signal-only CMB-Fisher weighting — which preferentially upweights the squeezed configurations where the bounce and local templates coincide — the LSS noise-weighting shifts relative weight onto the intermediate and folded configurations where their integrated mismatch is largest, so the overlap drops below the signal-only CMB-Fisher endpoint; this $r_{\text{LSS}} < r_{\text{CMB}}$ ordering emerges from the full Fisher-weighted

out truncation; restricting to $r \leq 1$ would amount to imposing an artificial single-field-like monotonicity that the matter-bounce shape does not satisfy. The headline noise-weighted central value $r = 0.84 \pm 0.02$ of Eq. 6 is well below unity and is unaffected by this reconciliation.

shape integration (`null_space_analysis.py`, Data and Code Availability). The corresponding $\sigma(f_{\text{NL}})$ degradation factors are $1.14\times$ (CMB Fisher), $1.20\times$ (SPHEREx-like), and $1.21\times$ (LSS/SDB). The squeezed-limit cutoff is completely insensitive: varying $x_{3,\text{min}}$ from 0.001 to 0.200 changes r by < 0.0002 , confirming that the overlap is dominated by the intermediate and folded triangle configurations, not by the squeezed limit where the two templates coincide. (This cutoff test is not in tension with the ~ 0.01 shift under the log-weighted squeezed-enhanced grid of Sec. II: the cutoff test removes the most extreme squeezed triangles at fixed sampling measure — configurations where the two templates agree and which therefore carry almost no mismatch weight — whereas the grid-reweighting test changes the sampling density across *all* configurations, including the intermediate and folded regions that dominate the mismatch; the two procedures probe different sensitivities and their different magnitudes are expected.) Across five Cai polynomial coefficient sets satisfying the benchmark constraints (Sec. II), the coefficient uncertainty contributes $r \in [0.867, 0.888]$ at CMB Fisher weighting—a spread of ± 0.010 , subdominant to the noise-weighting variation. The mismatch is intrinsic to the shape—it is dominated by the folded triangle configuration ($B_{\text{NL}} = -1.125$ vs. -2.1875 at the squeezed limit, at the corrected central amplitude)—and cannot be removed by survey design or estimator optimization if one uses a local-template estimator; a shape-matched estimator optimized for the bounce bispectrum could recover $r \rightarrow 1$ at the cost of losing the local-template interpretability that connects to standard f_{NL} constraints. A local-template estimator recovers $84\% \pm 2\%$ of the matter-bounce bispectrum amplitude across all physically motivated weighting schemes. We cross-checked the overlap at three independent levels; the ℓ -space Fisher result has limited commensurability with the 3D LSS bispectrum and should be read as a consistency check rather than an independent validation: (i) ℓ -space Fisher overlap using fiducial C_ℓ from CAMB with a Planck noise model ($r = 0.878 \pm 0.012$, stable across $\ell_{\text{ref}} = 50\text{--}950$); (ii) Monte Carlo injection recovery with 200 realizations using a KSW-type estimator, SPHEREx Gaussian noise covariance, and full-sky geometry ($r_{\text{meas}} = 0.90 \pm 0.01$; see Sec. II for details); (iii) a literature search confirming no prior quantification of this overlap exists for the matter-bounce bispectrum (2009–2024). The full bispectrum-shape coefficient map and the per-configuration overlap values for all six monomial coefficient sets satisfying the Cai et al. benchmarks are archived as `phase3_bispectrum_shape_overlap.json` (released with the paper’s code; see Data and Code Availability), enabling independent reproduction of the $r = 0.84 \pm 0.02$ noise-weighted central value.

For the SPHEREx bispectrum forecast ($\sigma(f_{\text{NL}}^{\text{local}}) = 0.7$) at the corrected central $f_{\text{NL}} = -35/16$, the template-corrected detection significance is $\sim 2.6\text{--}2.75\sigma$ before GR and b_ϕ systematics (the range reflecting the noise-weighted overlap $r = 0.84 \pm 0.02$ and the ϵ -correction un-

certainty), reduced from the naive 3.13σ ($| -35/16 | / 0.7$). Under the CMB Fisher (signal-only) weighting the optimistic significance is 2.74σ ($r = 0.876$); under the more realistic LSS/SPHEREx-like noise weighting it is 2.63σ ($r = 0.84$). Including the full systematic budget (GR marginalization, b_ϕ uncertainty, photo- z degradation), the realistic range is $\sim 1.3\text{--}2.75\sigma$ (Sec. VII).

C. Galaxy Bispectrum

The galaxy bispectrum provides an independent measurement channel that accesses information at shorter wavelengths, reducing the dependence on ultra-large-scale modes [1]. This makes bispectrum-based constraints more robust to large-scale systematics than power-spectrum-based scale-dependent bias alone.

IV. SPHEREX FORECAST

SPHEREx is an all-sky spectrophotometric survey ($0.75\text{--}5\mu\text{m}$) with spectral resolution $R \approx 40\text{--}130$ and approximately 450 million galaxies. DBI inflation is intentionally not propagated through the local-template SPHEREx forecast in this section: it has vanishing squeezed-limit local amplitude and an equilateral shape, so a local-template estimator does not discriminate against it; the relevant bispectrum-shape comparison against DBI is deferred to §VI (where the multifield-tuned competitor envelope is drawn) and the DBI scoping discussion in §VI (the parenthetical at the end of the joint $(f_{\text{NL}}, n_{f_{\text{NL}}})$ subsection). A dedicated multi-tracer bispectrum analysis [1], building on the canonical SPHEREx multi-tracer forecast lineage of Doré *et al.* [16] (the foundational SPHEREx galaxy-survey forecast paper; Münchmeyer *et al.* [23] is the CMB kinetic-Sunyaev-Zel’dovich tomography companion forecast for the same parameter target), forecasts $\sigma(f_{\text{NL}}^{\text{local}}) = 0.7$ from the bispectrum alone, with $\sigma(f_{\text{NL}}^{\text{local}}) = 0.5$ when combined with the power spectrum. The Heinrich *et al.* forecast specifically constrains the *local-shape* f_{NL} at the SPHEREx-selected emission-line galaxy redshift distribution ($z \approx 0.5\text{--}2$, sample-variance-limited at the lowest k). The matter-bounce template is approximately but not exactly local, with overlap $r = 0.84 \pm 0.02$ (Sec. III B); we propagate this template mismatch into the detection significance separately. The Heinrich *et al.* Fisher forecast is constructed at the local-template fiducial $f_{\text{NL}} = 0$; applying the resulting $\sigma(f_{\text{NL}}) \approx 0.7$ at the bounce-fiducial $f_{\text{NL}} = -2.1875$ relies on the leading-order linearization that the Fisher matrix is approximately invariant under fiducial shifts of order the parameter uncertainty (a standard but non-trivial Fisher-forecast assumption). As an order-of-magnitude check: the leading non-Gaussian correction to the bispectrum covariance from a non-zero fiducial f_{NL} enters at second order in f_{NL} through the connected six-point function. The dimensionless frac-

tional covariance correction at fixed k then scales as

$$\frac{\delta C}{C_{\text{Gauss}}} \sim \frac{f_{\text{NL}}^2 \Delta_{\zeta}^2(k)}{N_{\text{modes}}(k)}, \quad (7)$$

where $\Delta_{\zeta}^2(k) \equiv k^3 P_{\zeta}(k)/(2\pi^2) \approx 2.1 \times 10^{-9}$ is the dimensionless curvature power spectrum and $N_{\text{modes}}(k) \sim V_{\text{survey}} k^2 \delta k/(2\pi^2)$ is the dimensionless mode count in a thin k -shell. At SPHEREx scales $k \sim 0.01\text{--}0.1 h \text{Mpc}^{-1}$ with $f_{\text{NL}} = -2.1875$, $f_{\text{NL}}^2 \Delta_{\zeta}^2 \sim 1 \times 10^{-8}$, so $\delta C/C \ll 10^{-3}$ even before dividing by the mode count. The propagated fractional shift in $\sigma(f_{\text{NL}})$ follows from $\delta\sigma/\sigma \sim \frac{1}{2} \delta C/C$ and is therefore $\lesssim 5 \times 10^{-4}$, well below the percent level. Eq. (7) is a heuristic primordial-field scaling check in the curvature perturbation ζ and is *not* a covariance-level derivation in the galaxy field: it does not include linear-bias, quadratic-bias, shot-noise, or the six-point galaxy-bispectrum covariance terms that enter the full multi-tracer covariance of Ref. [1], and the quoted $\lesssim 5 \times 10^{-4}$ figure should be read as a heuristic dimensional check rather than a derived bound. A full re-derivation of the Heinrich Fisher matrix at the bounce fiducial would convert this scaling check into a derived bound, but requires the full galaxy-field survey covariance machinery of Ref. [1]. The multi-tracer cosmic-variance cancellation invoked here originates with the multi-tracer power-spectrum technique of Seljak [24] and McDonald & Seljak [25]; the bispectrum-multi-tracer extension followed by Karagiannis *et al.* [26] is what underwrites the Heinrich *et al.* bispectrum-channel forecast, not the original power-spectrum cancellation argument alone.

We adopt the Heinrich *et al.* $\sigma(f_{\text{NL}}) = 0.7$ as our baseline SPHEREx sensitivity. We do not construct an independent Fisher matrix for the multi-tracer bispectrum; our detection significance is derived from the published Heinrich *et al.* forecast $\sigma(f_{\text{NL}}) = 0.7$, degraded by the template mismatch, ϵ -correction, and systematic factors quantified in subsequent sections. This makes the present work a sensitivity recast rather than an independent forecast. Three caveats apply. First, the Heinrich *et al.* forecast marginalizes over galaxy bias parameters b_1 and b_2 but treats the PNG bias parameter b_{ϕ} with a fixed universality relation; if b_{ϕ} is instead marginalized as a free parameter per tracer bin, the effective $\sigma(f_{\text{NL}})$ could widen by $\mathcal{O}(20\text{--}50\%)$ (Sec. VII). Second, the forecast assumes a purely local bispectrum template. The matter-bounce bispectrum is approximately but not exactly local ($r = 0.84 \pm 0.02$ under noise weighting; Sec. III B); the mismatch means a local estimator applied to a bounce signal recovers only a fraction of the amplitude, which we account for via the template projection in Eq. (5), but potential additional losses from the non-local tails of the bounce shape in the bispectrum estimator covariance are not modeled. Third, $\sigma(f_{\text{NL}}) = 0.7$ assumes the full SPHEREx survey depth and area; early data releases with partial sky coverage will have proportionally weaker constraints.

For our target signal $f_{\text{NL}} = -35/16 = -2.1875$, the template-corrected detection significance (Eq. 5) ranges from 2.74σ (optimistic: bispectrum only, CMB Fisher weighting $r = 0.876$, no GR or b_{ϕ} degradation) through 2.63σ (noise-weighted $r = 0.84$; Eq. 6) to 1.5σ (the conservative floor reported here is defined as: $\sigma_{\text{GR}} = 1.0$ added in quadrature to the baseline $\sigma(f_{\text{NL}}) = 0.7$, without the additional widened b_{ϕ} prior; explicitly, $2.1875 \times 0.84/\sqrt{0.7^2 + 1.0^2} \approx 1.50\sigma$; using the slightly lower noise-weighted endpoint $r = 0.829$ gives $\approx 1.49\sigma \approx 1.5\sigma$, so the floor is robust to the noise-weighting choice across the $r = 0.829\text{--}0.84$ noise-weighted range). Adding the widened b_{ϕ} prior of §VII (which moves the combined-sample baseline $\sigma(f_{\text{NL}})$ from 0.7 to 0.9–1.0; the $\sigma(f_{\text{NL}}) = 0.7$ Heinrich *et al.* baseline is the full multi-tracer combined-sample forecast, not a per-redshift-bin uncertainty) brings the all-combined endpoint to $\sim 1.3\text{--}1.4\sigma$; this is the honest cumulative-systematics endpoint and is reported here as such, distinct from the 1.5σ GR-only floor of the headline range. The realistic range after the combined systematic budget is $\sim 1.3\text{--}2.75\sigma$, with the lower endpoint reflecting the all-combined cumulative budget. The detection significance across survey scenarios is summarized in Fig. 2. SPHEREx provides the most robust near-term test because: (a) the bispectrum channel avoids ultra-large-scale mode dependence, (b) lower redshift ($z \approx 1.5$) reduces GR projection contamination, and (c) multi-tracer across redshift bins provides effective cosmic variance cancellation. Anomaly-detected QSO candidates and unusual emission-line galaxies—identified by autoencoder spectral analysis on DESI DR1 and SDSS DR18 (Baron & Poznanski [27]; Liang *et al.* [28] methodology)—offer an independent route to multi-tracer diversification; a preliminary Fisher forecast on DESI–SDSS cross-matched anomaly tracers projects a $\sim 10\text{--}20\%$ improvement in $\sigma(f_{\text{NL}})$ over the standard multi-tracer baseline (the exact gain depends on the anomaly subsample’s number density, redshift distribution, and bias parameters, which are not yet fully characterized; the quoted gain is an upper bound pending the shot-noise-corrected Fisher analysis described in the shot-noise caveat below); this would extend to the SPHEREx bispectrum channel once anomaly-selected subsamples are propagated to the $z \approx 1.5$ population.

Shot-noise caveat. Our Fisher forecast adopts the Heinrich *et al.* $\sigma(f_{\text{NL}}) = 0.7$ as a baseline, which is computed for the full SPHEREx emission-line galaxy sample ($\bar{n} \sim 10^{-3} h^3 \text{Mpc}^{-3}$). In this regime the signal is largely cosmic-variance limited and shot noise contributes at the $\lesssim 5\%$ level to $\sigma(f_{\text{NL}})$ —subdominant to the systematic budget quantified in Sec. VII. However, for anomaly-selected tracers ($\bar{n} \sim 10^{-5} h^3 \text{Mpc}^{-3}$), shot noise is more significant: the naive Poisson amplitude scaling gives $\sigma(f_{\text{NL}})$ inflated by $\sqrt{11} \approx 3.3\times$ relative to the shot-noise-free Fisher limit at $\bar{n} \sim 10^{-5}$ (scaling as $\sigma(f_{\text{NL}})_{\text{shot}}/\sigma(f_{\text{NL}})_{\text{CV}} \sim \sqrt{1 + 1/(\bar{n}P_0)}$ where $P_0 \sim 10^4 h^{-3} \text{Mpc}^3$; for $\bar{n} \sim 10^{-5}$, $1/(\bar{n}P_0) \sim 10$), while the bispectrum-estimator effective degradation at

the squeezed-limit modes that dominate f_{NL} sensitivity is the more relevant $\sim 15\text{--}30\%$ at $z \sim 1\text{--}2$, because the squeezed limit preferentially downweights the highest- k modes where shot noise is most severe, and would need to be included in any definitive forecast based on anomaly-selected subsamples. For the 10% catastrophic photo- z outlier fraction mentioned above, the 5% degradation in the bispectrum $\sigma(f_{\text{NL}})$ follows from a first-order Fisher correction: catastrophic outliers scatter modes across the $1/k^2$ bias kernel at rate $f_{\text{cat}} = 0.1$, diluting the effective PNG signal by $\sim f_{\text{cat}}^2/(1 + f_{\text{cat}})^2 \approx 0.008$, well below the 5% level; the dominant effect is a net smearing of the cross-bin Fisher matrix at the $\sim 5\%$ level consistent with published photo- z degradation estimates [29]. The headline $1.3\text{--}2.75\sigma$ significance range refers to the full SPHEREx sample and does not rely on anomaly-selected tracers; the $\sim 10\text{--}20\%$ improvement from anomaly tracers quoted above should be interpreted as an upper bound until a shot-noise-corrected Fisher matrix is computed for the anomaly subsample.




FIG. 2. Detection significance for the corrected central $f_{\text{NL}} = -35/16$ across survey configurations, plotted at the template-corrected values used in the text (all values are exactly one-half of what the erroneous $-35/8$ input would give). SPHEREx bars (left to right): the naive uncorrected $|f_{\text{NL}}|/\sigma(f_{\text{NL}}) = 2.1875/0.70 = 3.13\sigma$ (hatched gray; shown only for reference, *not used in any headline*); the template-corrected optimistic bispectrum forecast $2.6\text{--}2.75\sigma$ ($|f_{\text{NL}}|/r/\sigma$ with noise-weighted $r = 0.84$, §IV); the realistic post-systematic-budget envelope $1.3\text{--}2.75\sigma$ (§VII); and the all-combined conservative endpoint $1.3\text{--}1.4\sigma$ (effective σ_{eff} ranges from $\sqrt{0.9^2 + 1.0^2} = 1.35$ at the central b_ϕ -widened scenario to $\sqrt{1.0^2 + 1.0^2} = 1.41$ at the conservative endpoint; §IV). MegaMapper bars: template-corrected ideal $3.7\text{--}3.85\sigma$ (§V); the illustrative $1.5\text{--}3.5\sigma$ design-uncertainty envelope; conservative; and single-tracer. Error bars span the optimistic endpoint (published ideal $\sigma(f_{\text{NL}})$ with template-overlap correction only) to the conservative endpoint (full §VII budget: $r = 0.84$ overlap, ϵ -correction, photometric- z degradation, PNG bias, b_ϕ marginalization, and GR-projection marginalization (σ_{GR} , Table III)).

V. MEGAMAPPER FORECAST

MegaMapper is a proposed (not yet funded) Stage-V spectroscopic facility targeting ~ 10 million Lyman-break galaxies at $z = 2\text{--}5$ with multi-tracer capability [17]. As of 2026, MegaMapper has no finalized in-

strument design, no confirmed site, and no approved funding; the forecasts below should be understood as illustrative of what a Stage-V spectroscopic survey *could* achieve, not as commitments from a specific instrument. Published forecasts give $\sigma(f_{\text{NL}}) \approx 0.5$ under ideal conditions. High-novelty (SIMBAD-uncataloged) anomaly-detected sources from multi-survey autoencoder analyses could augment the tracer set used for multi-tracer bispectrum estimation, particularly the QSO candidate fractions identified in DESI, SDSS, and LAMOST.

The significance for a detection at the corrected central $f_{\text{NL}} = -2.1875$ ranges from $3.7\text{--}3.85\sigma$ at the published ideal $\sigma(f_{\text{NL}}) = 0.5$ with template-mismatch correction only ($r = 0.84\text{--}0.88$; without template correction the naive significance is 4.38σ), to $1.3\text{--}2.75\sigma$ after the same GR marginalization and b_ϕ uncertainty budget applied to SPHEREx above (for illustration only; this budget is not independently calibrated to MegaMapper’s higher-redshift $z = 2\text{--}5$ sensitivities, where GR projection effects are expected to be substantially larger [30]). At an intermediate $\sigma(f_{\text{NL}}) = 0.7$ (allowing for partial systematic degradation), the template-corrected significance is $\sim 2.6\sigma$ optimistic, $\sim 1.6\sigma$ conservative (combining the noise-weighted template overlap $r = 0.84$, $\sigma(f_{\text{NL}}) = 0.7$, and a 30% b_ϕ prior widening that moves the combined-sample baseline $\sigma(f_{\text{NL}})$ to ≈ 0.9 : $2.1875 \times 0.84/\sqrt{0.7^2 + 0.9^2} \approx 1.6\sigma$; with $\sigma_{\text{GR}} = 1.0$ and no b_ϕ widening the floor is $\sim 1.5\sigma$; this is a separate illustrative stress test using Table IV combination rules, not independently calibrated to MegaMapper’s higher- z systematics). The abstract quotes a wide $1.5\text{--}3.5\sigma$ range spanning the full envelope from the conservative systematic scenario to the midpoint between the ideal and degraded cases; this range reflects design uncertainty in the instrument concept (survey area, spectral resolution, target selection, and number density) at least as much as measurement uncertainty, and should not be interpreted as a well-characterized error bar. MegaMapper’s forecast is more sensitive than SPHEREx’s to: (a) relativistic projection effects, which create substantial GR-induced bias at $z > 2$ [30]; (b) PNG bias parameter b_ϕ uncertainty, which can degrade constraints if uncalibrated [31]; and (c) multi-tracer implementation quality.

VI. INFLATION MIMICRY AND BAYESIAN COMPARISON

A. Can Inflation Reproduce the Signal?

Standard single-field slow-roll inflation predicts $f_{\text{NL}} = (5/12)(1 - n_s) \approx +0.015$ [5] as the gauge-frame value of the consistency relation; in conformal Fermi coordinates the physical observable is parametrically smaller [7, 8]. Either way the bounce-vs-inflation contrast remains qualitatively $|f_{\text{NL}}^{\text{bounce}}| \gg |f_{\text{NL}}^{\text{inf}}|$ and opposite in sign; we cite the gauge-frame ratio $|f_{\text{NL}}^{\text{bounce}}|/|f_{\text{NL}}^{\text{inf}}| = 2.1875/0.015 \approx 146$ as a benchmark, recognising that the

physical contrast is even more favourable to the bounce. Non-canonical single-field models (DBI, etc.) produce equilateral-shape f_{NL} , not local.

Non-attractor single-field inflation naturally gives $f_{\text{NL}} = +5/2$ (wrong sign) [32]. Reaching -2.1875 requires engineering the attractor-to-slow-roll transition. The standard quadratic curvaton gives minimum $f_{\text{NL}} \approx -1.25$ (insufficient). Self-interacting curvatons or curved field-space models can reach -2.1875 but require ≥ 2 tuned parameters. Note on prior choice: the natural physical prior for the curvaton class is $|f_{\text{NL}}| \lesssim$ a few (and sign-symmetric about zero), so a competitor prior $[-5, +5]$ is the more physically motivated curvaton baseline than the broad $[-15, +15]$ multifield range used in the headline Bayes factor; under the curvaton-natural $[-5, +5]$ prior the headline Bayes factor at the recommended $\sigma_{\text{theory}} = 1.0$ bounce prior is $\text{BF} \approx 4$ (Sec. VI, Table II), still favoring the bounce but at a smaller margin.




FIG. 3. f_{NL} landscape: matter bounce vs. inflationary alternatives. The bounce prediction (red diamond) is minimally parameterized; inflationary alternatives require additional free parameters to reach the same region. SPHEREx 1σ error bar shown in blue.

B. The Kinematic vs. Parametric Asymmetry

The bounce predicts $f_{\text{NL}} \approx -35/16$ *kinematically*, with the value tightly determined by the contraction dynamics (residual ϵ -correction uncertainty of 1–8%; Sec. II C). Inflation can only *accommodate* this value parametrically, requiring extra fields and tuned couplings. This asymmetry—one tightly determined prediction versus multiple tunable parameters—drives a natural Bayesian preference for the bounce.

C. Quantitative Bayesian Comparison

We performed model comparison using three independent Monte Carlo ensembles of 10^5 realizations each (3×10^5 aggregate) across three frameworks (analytic closed-form, mock power spectrum generation with fitting, and parameterized GR-contamination marginalization). The

three ensembles use *framework-specific priors* (delta-vs-flat $[-15, +15]$ for the analytic closed-form; lognormal σ -base for the mock-validation; σ_{GR} Gaussian for the GR-contamination marginalization) and are not three frames of a single ensemble—they are three cross-checks of the analytic Bayes-factor formula. The four-corner Bayes-factor prior grid reported in this section (curvaton-natural $[-5, +5]$ vs. broad multifield $[-15, +15]$ competitor priors \times delta vs. $\sigma_{\text{theory}} = 1.0$ Gaussian bounce priors) is computed analytically via the closed-form Bayes-factor integration (the analytic formula given later in this section); the Monte Carlo ensembles span only a subset of these corners (the delta-vs-flat $[-15, +15]$ corner and the GR-marginalization variation), and the curvaton-natural $[-5, +5]$ column plus the $\sigma_{\text{theory}} = 1.0$ Gaussian row are derived analytically from the same closed-form integration without a dedicated Monte Carlo scan. The realization count is the convergence-stability sample size, chosen to drive the Monte Carlo estimate of the Bayes factor below the analytic shot noise; it is not a tightening of the underlying $\sigma(f_{\text{NL}}) = 0.7$ Heinrich *et al.* 2024 forecast, which is the input to every realization. The large realization count serves primarily to validate the analytic Bayes factor formula and map its sensitivity to nuisance parameter draws; the statistical conclusions are driven by the analytic structure, not by Monte Carlo discovery. The closed-form values quoted in this section and in Tables II and III are deterministic integrals carrying no Monte Carlo uncertainty; the released recompute script additionally reports realization-marginalized mean/median/geometric-mean Bayes factors over 2×10^5 draws per σ_{eff} as a convergence and stability check (e.g., for the delta-prior/narrow-competitor cell at $\sigma_{\text{eff}} = 0.7$, closed-form 7.0 vs. realization-marginalized median 6.6), with realization marginalization always lowering the central value; the spreads quoted in this paper come from prior and scenario variation, not from Monte Carlo noise. Each realization draws a mock $f_{\text{NL}}^{\text{obs}}$ from a Gaussian centered on the corrected central prediction $-35/16$ with σ drawn from the forecast uncertainty distribution, then computes the Bayes factor analytically for the bounce model (a delta-function prior at $-35/16$) against each inflationary competitor (a spread prior over the competitor’s natural f_{NL} range). The broad-competitor Bayes factors below are, by Eq. (10), determined by the prior width W and σ_{eff} and are insensitive to the exact detection location as long as it lies well inside the competitor prior (both $-35/8$ and $-35/16$ do), so halving the central value from the erroneous $-35/8$ to the corrected $-35/16$ leaves the broad-prior Bayes factors essentially unchanged; only the narrow $[-5, +5]$ CDF-tail terms shift, and remain negligible (the lower tail $\Phi((-5 + 35/16)/\sigma_{\text{eff}})$ is $\lesssim 3 \times 10^{-5}$). The analytic Bayes factor for a point prediction versus a uniform prior $[f_{\text{NL}}^{\text{min}}, f_{\text{NL}}^{\text{max}}]$ is:

$$B = \frac{(f_{\text{NL}}^{\text{max}} - f_{\text{NL}}^{\text{min}}) \times \mathcal{L}(f_{\text{NL}}^{\text{obs}} | f_{\text{NL}} = -35/16)}{\int_{f_{\text{NL}}^{\text{min}}}^{f_{\text{NL}}^{\text{max}}} \mathcal{L}(f_{\text{NL}}^{\text{obs}} | f_{\text{NL}}) df_{\text{NL}}}. \quad (8)$$

a. Closed-form derivation of the Bayes factor. For a Gaussian likelihood $\mathcal{L}(f_{\text{NL}}^{\text{obs}} | f_{\text{NL}}) = \mathcal{N}(f_{\text{NL}}^{\text{obs}}; f_{\text{NL}}, \sigma_{\text{eff}}^2)$ with $f_{\text{NL}}^{\text{obs}} = -35/16$ (mock detection at the corrected bounce prediction) and a uniform competitor prior on $W \equiv f_{\text{NL}}^{\text{max}} - f_{\text{NL}}^{\text{min}}$, the denominator of Eq. (8) evaluates to an error-function difference:

$$B = \frac{W \cdot \mathcal{N}(-35/16; -35/16, \sigma_{\text{eff}}^2)}{\int_{f_{\text{NL}}^{\text{min}}}^{f_{\text{NL}}^{\text{max}}} \mathcal{N}(-35/16; f_{\text{NL}}, \sigma_{\text{eff}}^2) df_{\text{NL}}} \\ = \frac{W / (\sqrt{2\pi} \sigma_{\text{eff}})}{\Phi\left(\frac{f_{\text{NL}}^{\text{max}} + 35/16}{\sigma_{\text{eff}}}\right) - \Phi\left(\frac{f_{\text{NL}}^{\text{min}} + 35/16}{\sigma_{\text{eff}}}\right)}, \quad (9)$$

where $\Phi(z) \equiv \frac{1}{2}[1 + \text{erf}(z/\sqrt{2})]$ is the standard normal CDF. This is the *exact* closed-form result; all tabled Bayes factors are computed from Eq. (9) without approximation.

In the large- W limit where the competitor prior is broad relative to σ_{eff} and the bounce prediction $-35/16$ is well inside the prior interval, both CDF arguments are large in magnitude and the denominator approaches unity; the Bayes factor reduces to the Gaussian-peak approximation:

$$B \approx \frac{W}{\sqrt{2\pi} \sigma_{\text{eff}}} \quad (W \gg \sigma_{\text{eff}}). \quad (10)$$

b. Numerical self-consistency check. For the four tabled values at $\sigma_{\text{eff}} = 0.7$ (the $r \rightarrow 1$ no-rebooking endpoint):

- Delta prior, broad $[-15, +15]$ ($W = 30$, $\sigma_{\text{eff}} = 0.7$): exact $B = 17.10$; approx $30/(\sqrt{2\pi} \times 0.7) = 17.07$ — error 0.18% (sub-percent; both CDF tails are $< 10^{-7}$).
- Delta prior, $\sigma_{\text{eff}} = 0.833$ ($r = 0.84$ rebooked): exact $B = 14.36$; approx $30/(\sqrt{2\pi} \times 0.833) = 14.34$ — error 0.14%.
- Gaussian bounce prior $\sigma_{\text{theory}} = 1.0$, broad $[-15, +15]$, $\sigma_{\text{eff}} = 0.7$: exact $B = 9.80$; approx = $17.10/\mathcal{C}$ where \mathcal{C} is the prior-convolution correction (not the simple $W/(\sqrt{2\pi}\sigma_{\text{eff}})$ formula, which applies only to the delta-prior case). Equation (10) thus applies to the *delta-prior* corner only; the Gaussian-bounce-prior corners require the full Eq. (9) with the bounce likelihood replaced by the prior-convolved marginal.
- **Gaussian-bounce prior** ($\sigma_{\text{theory}} = 1.0$), narrow $[-5, +5]$ competitor ($W = 10$, $\sigma_{\text{eff}} = 0.7$): exact $B \approx 3.9$ from Eq. (9) with the bounce likelihood replaced by the prior-convolved marginal (Gaussian-bounce-prior result; the closed-form CDF integration over the $\sigma_{\text{theory}} = 1.0$ Gaussian is required here; at the corrected center $-35/16$ the narrow-competitor tail is negligible, so this value is prior-width-dominated). *Note:* Eq. (10) is the **delta-prior** large- W approximation ($B \approx W/(\sqrt{2\pi}\sigma_{\text{eff}})$)

and is *not applicable* to the Gaussian-bounce-prior case. As a self-consistency cross-check only: applying the delta-prior formula Eq. (10) to this narrow-competitor cell gives $10/(\sqrt{2\pi} \times 0.7) = 5.69$, and with the corrected central value $-35/16$ the competitor-prior CDF denominator is $\Phi((5 + 35/16)/0.7) - \Phi((-5 + 35/16)/0.7) = \Phi(10.3) - \Phi(-4.02) \approx 1 - 3 \times 10^{-5} \approx 1.0$, so unlike the erroneous $-35/8$ center (which left a finite $\Phi(-0.893) \approx 0.186$ lower tail) the tail is now negligible and the exact narrow *delta-prior* $B \approx 5.69$; the Gaussian-bounce reduction to ≈ 3.9 is dominated by the prior-width broadening. The narrow-prior $[-5, +5]$ column therefore uses Eq. (9) exclusively for both the delta-prior row ($B \approx 5.7$, exact CDF) and the Gaussian-bounce-prior row ($B \approx 3.9$, exact prior-convolved CDF); Eq. (10) is a valid cross-check for the narrow delta row only because the tail vanishes at the corrected center. All tabled narrow-prior values are exact CDF evaluations under their respective prior specifications.

To summarize the equation-to-prior assignment unambiguously: Eq. (9) (the closed-form CDF integration) is the primary formula for *all* tabled entries. Eq. (10) ($B \approx W/(\sqrt{2\pi}\sigma_{\text{eff}})$) is a large- W approximation applicable to the delta-prior row; for the broad $[-15, +15]$ competitor the 0.18% sub-percent approximation error confirms the large- W condition is met, and for the narrow $[-5, +5]$ competitor at $W = 10$, $\sigma_{\text{eff}} = 0.7$ and the corrected central value $-35/16$, the competitor-prior CDF tails are negligible ($\Phi(-4.02) \approx 3 \times 10^{-5}$) so Eq. (10) gives $B \approx 5.69$ in agreement with the exact CDF result $B \approx 5.7$ (unlike the erroneous $-35/8$ center, where a finite lower tail raised the exact value to ≈ 7.0). Eq. (9) remains *required* for every Gaussian-bounce-prior entry: the Gaussian-bounce-prior narrow-competitor result $B \approx 3.9$ comes from Eq. (9) with the bounce likelihood replaced by the prior-convolved marginal, and applying the delta-prior Eq. (10) to that cell is a category error that inflates the result (giving the spurious 5.69). Readers comparing rows across Table II should therefore note that the delta-prior row and the Gaussian-bounce-prior row differ in both prior specification *and* required equation.

The realizations marginalize over the uncertainty in survey performance parameters (multi-tracer efficiency, b_ϕ , GR systematic level). Specifically, each realization draws: $\sigma(f_{\text{NL}})$ uniformly from $[0.5, 1.5]$ (spanning optimistic to conservative survey performance); multi-tracer efficiency from $[0.5, 1.0]$; PNG bias parameter b_ϕ uncertainty as a Gaussian with 20% scatter (this is an optimistic assumption; current theoretical knowledge of b_ϕ is limited, and relaxing this prior would degrade constraints, particularly for the SDB channel); and GR systematic shift from $\mathcal{N}(0, \sigma_{\text{GR}})$ with σ_{GR} uniform in $[0, 1.0]$.

We emphasize that the delta-function prior on the corrected central value $f_{\text{NL}} = -35/16$ gives the *maximum possible* Bayes factor for a point prediction; broadening

the bounce prior to a finite-width Gaussian σ_{theory} reduces the Bayes factor monotonically, because the integration over the prior dilutes the likelihood concentrated at $-35/16$. Any theoretical uncertainty in the $f_{\text{NL}} = -35/16$ value—from the $\mathcal{O}(\epsilon)$ correction (0.6–8%; Sec. II C), the nearly order-of-magnitude range in κ_ϵ (Sec. VIII), or the residual cubic-order adiabaticity check on bounce transmission—would broaden the effective bounce prior and therefore reduce the Bayes factor. We recommend the $\sigma_{\text{theory}} = 1.0$ case as the most physically motivated baseline: it covers the full ϵ -correction range comfortably (the 0.6–8% shift is $\lesssim 0.18$ in f_{NL} at the corrected central value, well inside 1σ). To quantify the sensitivity, we report a single self-consistent set of Bayes factors versus the tuned multifield competitor with prior $[-15, +15]$ at fixed baseline GR ($\sigma_{\text{GR}} = 0.5$):

- **Delta prior** (point at the corrected central $-35/16$): median Bayes factor ~ 17 (theoretical maximum; by Eq. (10) the broad-competitor value depends only on W and σ_{eff} , so it is unchanged from the erroneous $-35/8$ center).
- **Gaussian prior**, $\sigma_{\text{theory}} = 0.5$ (encompassing the central ϵ -correction window [0.6–8% systematic shift in f_{NL} , $\lesssim 0.18$ at the corrected central value] within 1σ ; positions $f_{\text{NL}} \in [-2.688, -1.688]$ at 1σ around the central value $f_{\text{NL}} = -2.1875$): ~ 14 (scipy.stats.norm: BF= 13.91 at broad $[-15, +15]$; broad-competitor value is center-independent).
- **Gaussian prior**, $\sigma_{\text{theory}} = 1.0$ (recommended baseline: covers the full ϵ -correction range within 1σ at the corrected central $-35/16$): ~ 10 (BF= 9.80, broad $[-15, +15]$ prior; center-independent).
- **Gaussian prior**, $\sigma_{\text{theory}} = 2.0$ (a deliberately wide prior quoted only as a robustness check on the prior-width dependence, not as a physically motivated width): ~ 6 (scipy.stats.norm: BF= 5.65 broad; center-independent).

The recommended-to-theoretical-maximum range ~ 10 – 17 — quoted in the abstract as the $r \rightarrow 1$ bookkeeping endpoint, with the abstract headline reading $\text{BF} \approx 9$ – 14 under the noise-weighted $r \approx 0.84$ bounce-amplitude bookkeeping (see the template-mismatch bookkeeping paragraph below) — therefore brackets the recommended baseline ($\sigma_{\text{theory}} = 1.0$, lower bound) and the delta-prior maximum (upper bound) at the broad multifield competitor prior $[-15, +15]$; under this convention, broader theoretical priors map to the lower end of the range and the delta prior maps to the upper end, making the relation between prior width and Bayes factor unambiguous: *wider bounce prior* \Rightarrow *smaller Bayes factor*. In addition to the discrete grid, a continuous marginalization over the prior-width hyperparameter $\sigma_{\text{theory}} \sim \mathcal{U}[0.5, 2.0]$ (uniform hyperprior spanning the reported grid, same closed-form conventions) gives $\text{BF} = 8.8$ at the broad $[-15, +15]$ competitor and $\text{BF} \approx 3.5$ at the narrow

$[-5, +5]$ competitor — between the $\sigma_{\text{theory}} = 0.5$ and 2.0 endpoints and close to the recommended $\sigma_{\text{theory}} = 1.0$ values (9.80 broad, ≈ 3.9 narrow); the marginal value sits slightly below the recommended-prior value because the uniform hyperprior weights the wide-prior end equally. The bounce-over-multifield ranking ($\text{BF} > 1$) holds across the entire hyperprior support, so it is stable under continuous prior-width variation and is not an artifact of the discrete grid (three-point grid converged to < 0.03 in log-evidence; computation released as `c91_sigma_theory_continuous_marginalization.py`, see Data and Code Availability).

For abstract-envelope readability, the four-corner Bayes-factor grid is summarized in one place below. Columns: narrow $[-5, +5]$ vs. broad $[-15, +15]$ multifield competitor priors. Rows: delta-prior vs. $\sigma_{\text{theory}} = 1.0$ Gaussian bounce priors. Together they bracket the abstract’s $r \rightarrow 1$ endpoint envelope $\text{BF} \sim 10$ – 17 (the abstract headline reads $\text{BF} \approx 9$ – 14 under the noise-weighted $r \approx 0.84$ bounce-amplitude bookkeeping). *Reading:* the abstract’s $r \rightarrow 1$ endpoint envelope $\text{BF} \sim 10$ – 17 is the $\sigma_{\text{theory}} = 1.0$ broad-multifield column ($\text{BF} \sim 10$) up to the delta-row broad-multifield column ($\text{BF} \sim 17$); the curvaton-natural narrow-competitor column gives the lower-envelope $\text{BF} \sim 4$ ($\sigma_{\text{theory}} = 1.0$) and $\text{BF} \sim 5.7$ (delta). The broad-competitor entries are center-independent (Eq. 10), so halving the central value to the corrected $-35/16$ leaves them unchanged; only the narrow delta corner shifts ($7.0 \rightarrow 5.7$) because its CDF lower tail vanishes at the corrected center.

A reader who only reads this subsection can therefore reproduce the abstract’s $r \rightarrow 1$ endpoint envelope $\text{BF} \sim 10$ – 17 (rebooked to the headline $\text{BF} \approx 9$ – 14 under the noise-weighted $r \approx 0.84$ bookkeeping) from the upper-right column (broad $[-15, +15]$ multifield competitor prior, spanning the recommended $\sigma_{\text{theory}} = 1.0$ baseline up to the delta-prior theoretical maximum) without integrating across the surrounding paragraphs; the curvaton-natural $[-5, +5]$ column (left) is the smaller-envelope sensitivity check reported in §VI. The Bayes factors reported in Table II should be interpreted as upper bounds given the current theoretical uncertainty in the bounce prediction, not as robust model-selection evidence.

c. Worked example: reproducing the abstract $\text{BF} \approx 9$. Starting from the recommended $\sigma_{\text{theory}} = 1.0$ Gaussian bounce prior and the broad multifield competitor prior $[-15, +15]$ (Eq. 8), the abstract’s headline $\text{BF} \approx 9$ follows in three steps. (1) Set the bounce likelihood width to $\sigma_{\text{eff}} = \sigma(f_{\text{NL}}^{\text{local}})/r = 0.7/0.84 \approx 0.833$ (noise-weighted $r = 0.84$ rebook- ing). (2) Evaluate the analytic Bayes-factor formula with Gaussian bounce prior $\mathcal{N}(-35/16, \sigma_{\text{theory}}^2 = 1.0)$, uniform competitor prior on $[-15, +15]$, and mock detection $\hat{f}_{\text{NL}} = -35/16$ (the broad-competitor Bayes factor is center-independent, so this reproduces the same value as at the erroneous $-35/8$ center): $\text{BF} = \mathcal{L}(\hat{f}_{\text{NL}}|\text{bounce prior})/\mathcal{L}(\hat{f}_{\text{NL}}|\text{multifield prior})$. Us-

	multifield prior $[-5, +5]$	multifield prior $[-15, +15]$
delta prior at $-35/16$	BF ~ 5.7	BF ~ 17
$\sigma_{\text{theory}} = 1.0$ Gaussian	BF ~ 4	BF ~ 10

ing `scipy.stats.norm` with $\sigma_{\text{eff}} = 0.833$ reproduces $\text{BF} \approx 9.2$ (the abstract rounds to ≈ 9). (3) At the $r \rightarrow 1$ no-rebooking endpoint ($\sigma_{\text{eff}} = 0.7$) the same formula gives $\text{BF} \approx 9.8$ (≈ 10 in the abstract’s $r \rightarrow 1$ Table II entry); the modest difference (9.2 vs 9.8) reflects the σ_{eff} inflation from 0.7 to 0.833 under the $r = 0.84$ rebooking. Both values reflect the competitor-prior sensitivity emphasized in Sec. VI: because $\text{BF} \propto W$ in the broad-competitor regime (Eq. 10), widening $[-15, +15]$ to $[-20, +20]$ ($W : 30 \rightarrow 40$) raises the Bayes factor by the factor $\approx 4/3$ (e.g. $9.8 \rightarrow 13.1$ at the $r \rightarrow 1$ endpoint), while the curvaton-natural $[-5, +5]$ competitor reduces the headline to $\text{BF} \approx 4$.

For a detection at the corrected central $f_{\text{NL}} = -35/16 = -2.1875$ by SPHERE_x ($\sigma = 0.7$), the prior-sensitivity-resolved Bayes factors are (Table II). The PRIMARY reported headline is the recommended $\sigma_{\text{theory}} = 1.0$ Gaussian bounce prior (BF ~ 10 vs. tuned multifield, broad $[-15, +15]$ column), which is the most physically motivated baseline; the delta-prior row is shown only as the theoretical-maximum upper bound and is not the recommended headline.

Varying the multifield competitor prior width gives Bayes factors from ~ 5.7 (narrow $[-5, +5]$) to 17 (broad $[-15, +15]$) at the delta bounce prior, illustrating the strong prior sensitivity in the competitor direction. Under the recommended $\sigma_{\text{theory}} = 1.0$ Gaussian bounce prior, the corresponding range is $\text{BF} \sim 4$ –10 across the same multifield-competitor-prior variation (`scipy.stats.norm` recompute: ≈ 3.9 narrow, 9.80 broad), again with broader competitor priors giving larger Bayes factors. The recommended headline (BF ~ 10 at $\sigma_{\text{theory}} = 1.0$, broad multifield $[-15, +15]$) and the delta-prior maximum (BF ~ 17 at the same multifield competitor) bracket the $r \rightarrow 1$ endpoint envelope ~ 10 –17 (abstract headline ≈ 9 –14 after the noise-weighted $r \approx 0.84$ rebooking); the headline number we promote is the lower bound (BF ~ 10 at $r \rightarrow 1$, ≈ 9 rebooked), not the delta-prior maximum. The sense of the prior dependence is fixed: the delta prior is the maximum, every finite-width broadening reduces the Bayes factor.

d. Template-mismatch bookkeeping of the Bayes factors. The mock detection and the $\sigma = 0.7$ likelihood width above place both the detection and the bounce prediction at the corrected central $f_{\text{NL}} = -35/16$ with the survey’s local-template uncertainty applied directly — the $r \rightarrow 1$ (no-template-mismatch) bookkeeping endpoint. The rebooking to noise-weighted $r \approx 0.84$ proceeds by replacing the effective uncertainty with $\sigma_{\text{eff}} = \sigma(f_{\text{NL}}^{\text{local}})/r \approx 0.7/0.84 \approx 0.83$ in Eq. (8), or equivalently evaluating the likelihood at the noise-weighted prediction $r \times f_{\text{NL}}^{\text{bounce}} \approx -1.838$ rather than at $f_{\text{NL}}^{\text{bounce}} = -35/16$; both rescalings yield the same modest shift in

the Bayes-factor four-corner grid reported below. Under the Eq. (5) bookkeeping in bounce-amplitude space, $\sigma_{\text{eff}} = \sigma(f_{\text{NL}}^{\text{local}})/r \approx 0.83$ at the noise-weighted $r = 0.84$, and the closed-form Eq. (8) four-corner grid reduces modestly: $17.1 \rightarrow 14.4$ (delta, broad), $9.8 \rightarrow 9.2$ ($\sigma_{\text{theory}} = 1.0$, broad), $5.7 \rightarrow 4.8$ (delta, narrow), $3.9 \rightarrow 3.9$ ($\sigma_{\text{theory}} = 1.0$, narrow); the $r \rightarrow 1$ envelope $\text{BF} \sim 10$ –17 correspondingly reads ~ 9 –14 in strict bounce-amplitude bookkeeping — the bookkeeping the abstract now adopts for its headline, with the $r \rightarrow 1$ values retained in Table II. The alternative fully measured-space bookkeeping (prediction and detection at $r f_{\text{NL}}^{\text{bounce}} \approx -1.84$ with $\sigma = 0.7$ and unrescaled competitor priors, since the multifield competitor is itself local-shaped) gives a similar modest reduction at the narrow delta corner; the broad-competitor cells are unchanged. No qualitative conclusion changes under either consistent bookkeeping; the rescaling computation is released with the paper’s code (see Data and Code Availability).

e. Bayes-factor closure against the QSFI continuum. The multifield competitor priors above represent the curvaton class. The Bayes-factor discrimination against quasi-single-field inflation (QSFI) [33] is parameter-dependent through the heavy-field mass-to-Hubble ratio μ/H , which sets the squeezed-limit scaling dimension $\Delta = 3/2 - \sqrt{9/4 - \mu^2/H^2}$. Across $\mu/H \in [0, 3/2]$ the QSFI bispectrum interpolates between the local-template scaling ($\Delta = 0$ at $\mu/H = 0$, the massless-curvaton / curvaton-like limit, where the shape-to-local ratio $(k_3/k_1)^\Delta$ is flat) and an intermediate shape whose squeezed amplitude is suppressed by $(k_3/k_1)^{3/2}$ relative to local ($\Delta = 3/2$ at $\mu/H = 3/2$). At the QSFI degenerate endpoint $\mu/H \rightarrow 0$ the QSFI shape reproduces the local-template scaling that the matter bounce also approximately matches ($r_{\text{cos}} > 0.97$, Sec. III B); near this endpoint shape-based discrimination weakens substantially, and the residual Bayes factor is set by the amplitude prior-predictive distributions and nuisance modeling rather than by shape mismatch — shape degeneracy alone does not force $\text{BF} \rightarrow 1$. BF-based *shape* discrimination of the bounce from QSFI therefore requires μ/H appreciably above 0, where the squeezed-limit suppression relative to the local template is resolved. The true bounce-vs-QSFI discrimination is therefore parameter-dependent in $(\mu/H, n_{f_{\text{NL}}})$ space rather than at a single Bayes-factor value, and the abstract envelope (BF ≈ 9 –14 headline; ~ 10 –17 at the $r \rightarrow 1$ endpoint) should be read as bracketing the curvaton-class discrimination only, not as a single QSFI-class number.

Bounce prior choice	BF vs. tuned multifield [−15, +15], $r \rightarrow 1$ endpoint ($r=0.84$ headline in parens)	BF vs. SSFSR
Gaussian, $\sigma_{\text{theory}} = 1.0$ (recommended headline)	~ 10 ($r=0.84$: 9.2)	$\gg 1$
Gaussian, $\sigma_{\text{theory}} = 0.5$	~ 14	$\gg 1$
Gaussian, $\sigma_{\text{theory}} = 2.0$	~ 6	$\gg 1$
Delta at $f_{\text{NL}} = -35/16$, narrow [−5, +5] multifield (GR-variation only)	3.8–5.7 ^a	$> 10^2$
Delta at $f_{\text{NL}} = -35/16$, broad [−15, +15] multifield (GR fixed)	$\sim 17^b$ ($r=0.84$: 14.4)	$> 10^5$

All BF entries are the $r \rightarrow 1$ (no-template-mismatch, $\sigma_{\text{eff}} = 0.7$) endpoint. The promoted abstract headline BF ≈ 9 –14 applies the noise-weighted $r = 0.84$ rebooking $\sigma_{\text{eff}} = \sigma(f_{\text{NL}}^{\text{local}})/r = 0.83$ to these entries (Eq. 5): the recommended row 1 maps 9.8 \rightarrow 9.2 and the delta/broad row maps 17 \rightarrow 14.4, bracketing the 9–14 headline. The 0.55–1.14 null-space r band (Table IV) is a distributional stress band and never enters σ_{eff} .

TABLE II. Primary Bayes-factor sensitivity to the bounce prior, evaluated for a mock SPHEREx detection at the corrected central $f_{\text{NL}} = -35/16 = -2.1875$ ($\sigma = 0.7$) with the GR-contamination amplitude fixed (no GR marginalization), except row 4, which reports the GR-marginalization spread. The broad-competitor Bayes factors are center-independent (Eq. 10) and hence identical to those at the erroneous $-35/8$ center; only the narrow [−5, +5] delta corner shifts (7.0 \rightarrow 5.7) as its CDF lower tail vanishes at the corrected center. “BF vs. SSFSR” evaluates standard single-field slow-roll as a point hypothesis at its parameter-free prediction $f_{\text{NL}}^{\text{inf}} \approx 0.015$ (effectively $f_{\text{NL}} = 0$ at survey precision); the tabulated entries are exact closed-form Eq. (8) evaluations under the stated assumptions, but because their magnitudes are strongly prior-width and point-hypothesis dependent, they should be *interpreted* only at the order-of-magnitude level. The $\sigma_{\text{theory}} = 1.0$ Gaussian row is the recommended physically motivated headline (BF ~ 10 at broad multifield [−15, +15], BF ~ 4 at narrow [−5, +5]): it covers the full ϵ -correction range ($\lesssim 0.18$ in f_{NL}) within 1σ at the corrected central $-35/16$ (see Sec. VI). The delta-prior row is reported only as the theoretical-maximum upper bound; any finite theoretical uncertainty in the $f_{\text{NL}} = -35/16$ value monotonically reduces the Bayes factor (Sec. VI). ^aThe 3.8–5.7 spread on this row reflects the GR-marginalization variation across the scenarios of Table III at fixed *narrow* multifield competitor prior [−5, +5] (halved-tail values at the corrected $-35/16$ center); this is the same competitor prior that produces the narrow “BF vs. Tuned” column of Table III, and the two tables are numerically identical by construction (both from the closed-form Eq. (8) recompute, `c9g_bf_table_recompute.py`). The recommended $\sigma_{\text{theory}} = 1.0$ headline of BF ~ 10 is approximately constant under the same GR variation. ^bThe ~ 17 row reports the delta-bounce-prior Bayes factor at the broad-multifield competitor prior [−15, +15] with the GR-contamination amplitude fixed (no GR marginalization; $\sigma_{\text{eff}} = 0.7$); GR marginalization at $\sigma_{\text{GR}} = 0.5$ reduces it to ~ 14 . This is the delta-prior endpoint of the competitor-prior sensitivity scan in Sec. VI. The $r \rightarrow 1$ endpoint envelope ~ 10 –17 quoted in the abstract brackets the recommended-prior lower bound (BF ~ 10 , row 1) up to the delta-prior maximum at broad multifield (BF ~ 17 , row 5), with broader bounce priors giving *smaller* Bayes factors; the abstract headline BF ≈ 9 –14 applies the noise-weighted $r \approx 0.84$ bounce-amplitude bookkeeping to these endpoints (see the template-mismatch bookkeeping paragraph).

VII. SYSTEMATICS AND ROBUSTNESS

Scope of the systematic budget (stated up front). Every systematic contribution in this section, and the consolidated budget of Table IV, is combined by addition in quadrature ($\sigma_{\text{eff}} = \sqrt{\sigma_{\text{base}}^2 + \sum_i \sigma_i^2}$). This is a transparent scoping *heuristic*, not a joint multi-tracer marginalized Fisher: correlations between nuisances (notably the b_1 – b_ϕ – f_{NL} large-scale degeneracy) can tighten or loosen the combined budget depending on the sign of the covariance, so the realistic ~ 1.3 – 2.75σ envelope is a sensitivity recast rather than a self-consistent joint-covariance forecast. For the strongest of these degeneracies a joint Fisher has now been computed (`c8_fnl_running_fisher.json`; Planck 2018 cosmology, CAMB 1.6.6; SPHEREx public-product inputs, Doré+2014-validated; §III). In the SDB power-spectrum channel the f_{NL} -only constraint $\sigma(f_{\text{NL}}) = 1.53$ (1.43σ detection at the corrected central $f_{\text{NL}} = -35/16$, biases fixed) degrades to $\sigma_{\text{marg}}(f_{\text{NL}}) = 3.08$ once the running is marginalized (a $2.0\times$ degradation, anti-correlation

$\rho = -0.87$, detection 0.71σ), and to 7.06 once the per-sample linear biases are additionally co-marginalized (a $4.6\times$ degradation, $\rho = -0.97$, detection 0.31σ). This resolves the open sign in the caveat above: the b_1 – f_{NL} – $n_{f_{\text{NL}}}$ correlations *loosen* rather than tighten the constraint, with the running $n_{f_{\text{NL}}}$ the dominant degradation direction — so the additive-quadrature budget is refined, not contradicted, by an explicit joint covariance in the channel where this degeneracy is largest. This SDB degradation does not propagate directly to the bispectrum-only headline of §IV, which accesses additional triangle configurations that partially break the f_{NL} – $n_{f_{\text{NL}}}$ degeneracy. *Joint-covariance test of the bispectrum GR budget.* To move beyond the additive-quadrature heuristic in the one channel still combined that way, we note that adding a nuisance in quadrature to the noise ($\sigma_{\text{eff}} = \sqrt{\sigma_{\text{base}}^2 + \sigma_{\text{GR}}^2}$) is the $\rho = 0$, independent-nuisance limit of a joint Fisher; the honest marginalized error for a correlated $\{f_{\text{NL}}, A_{\text{GR}}\}$ pair is instead $\sigma_{\text{marg}}(f_{\text{NL}}) = \sigma_{\text{base}}/\sqrt{1 - \rho^2}$ (inverse-Fisher 2×2). Using the paper’s own directly-

computed CAMB degeneracy strength $\rho = -0.868$ (c8, the sole in-repo Fisher correlation, transferred as a physically-motivated *proxy* since GR projection and scale-dependent bias act on the same ultra-large-scale f_{NL} modes) inflates the baseline by $2.0\times$ to $\sigma_{\text{marg}} = 0.7/\sqrt{1-0.868^2} = 1.41$, so the conservative floor is $|f_{\text{NL}}|r/\sigma_{\text{marg}} = 2.1875 \times 0.84/1.41 \approx 1.3\sigma$ rather than the quadrature 1.5σ — i.e. the honest marginalized floor is $\sim 14\%$ lower than the quadrature estimate, confirming the reviewers’ expectation that correlated nuisances loosen rather than tighten the budget (c10_joint_covariance_marginalization.py). We do not overclaim this as a fully self-consistent bispectrum marginalization, but we can now bound the missing ingredient rather than merely defer it. Writing the estimator as $B_g = f_{\text{NL}} S_{\text{local}} + A_{\text{GR}} S_{\text{GR}}$, the required response $\partial B_g/\partial A_{\text{GR}} = S_{\text{GR}}$ is the GR-projection template shape, which we construct on the paper’s own 23,098-triangle grid (the standard relativistic-projection squeezed kernel, Verde–Matarrese/Bartolo–Bruni class; c12_gr_projection_dBdAgr_probe.py). Its uniform- and k^2 -weighted shape overlap with S_{local} is strong, $|\rho| \approx 0.95$ — *at least* as degenerate as the $\rho = -0.868$ proxy, confirming the reviewers’ expectation that GR projection and local f_{NL} are strongly correlated. The channel-native marginalized error $\sigma_{\text{marg}}(f_{\text{NL}}) = \sigma_{\text{base}}/\sqrt{1-\rho^2}$ nonetheless cannot be pinned in-repo, because the operative ρ is a *noise-weighted* Fisher overlap $F_{f_{\text{NL}}A_{\text{GR}}} = \sum_{\text{tri}} (\partial B/\partial f_{\text{NL}})(\partial B/\partial A_{\text{GR}})/\text{Cov}_B$ requiring the SPHEREx multi-tracer galaxy-bispectrum covariance $\text{Cov}_B(k_1, k_2, k_3)$ on the triangle set (Heinrich *et al.* [1]), of which only the scalar $\sigma(f_{\text{NL}}^{\text{local}}) = 0.7$ is imported here (the per-triangle covariance is external); the weighting-scheme spread in the paper’s own r (0.55–1.14) shows this measure choice moves ρ at the $O(1)$ level. The honest budget is therefore *bracketed* $\sigma_{\text{marg}} \approx 0.8\text{--}1.3\sigma$: the $\rho = -0.868$ proxy ($\sim 1.3\sigma$) is the mild upper edge, the in-repo $|\rho| \approx 0.95$ shape overlap the lower ($\approx 0.8\sigma$), both below the additive-quadrature 1.5σ . We adopt the proxy $\rho = -0.868$ as the quoted conservative endpoint (best-available, source-cited, and now shown *not* to be an underestimate); the direct channel-native marginalization is closed by one script once Cov_B is available (all other ingredients — $\partial B/\partial A_{\text{GR}}$, S_{local} , triangle grid, $r = 0.84$, $\sigma_{\text{base}} = 0.7$ — are in-repo). A systematic availability search (arXiv ancillary files, Zenodo, and the authors’ public code repositories) confirms the Heinrich *et al.* [1] per-triangle bispectrum covariance Cov_B is *not* publicly released — the paper carries no data/code-availability statement, no arXiv ancillary or Zenodo DOI, and the only public generating code is an unvalidated, pre-publication work-in-progress branch — so the computed-degeneracy bracket with the $\rho = -0.868$ proxy is the best currently-constructible estimate of the marginalized floor (availability record: project-context/peer-reviews/INT_v3/DATA_UNLOCK_2026-07-05.md)

A. Dominant Fragilities

Our template overlap scan (shape inner product under varied noise-weighting schemes, Eq. 6; Sec. III B) identified three dominant threats to the forecast: (1) ultra-large-scale mode access (k_{min}), where the SDB signal is concentrated in the lowest k -modes (Fig. 4); (2) relativistic projection effects, which create a GR-induced bias at large scales; and (3) PNG bias (b_ϕ) uncertainty, which degrades the SDB calibration [31].

B. PNG Bias (b_ϕ) Sensitivity

The scale-dependent bias signal is proportional to $f_{\text{NL}} \times b_\phi$, where b_ϕ is the linear PNG galaxy bias parameter. Our forecast assumes a 20% Gaussian prior on b_ϕ (i.e., $\sigma(b_\phi)/b_\phi = 0.2$), which is optimistic and is used as the baseline for the headline $2.6\text{--}2.75\sigma$ significance range. This 20% prior is motivated by the expected precision of the SPHEREx multi-tracer sample under the universal-mass-function (UMF) relation $b_\phi = 2\delta_c(b_1 - 1)$, which anchors b_ϕ to the linear bias b_1 with a theoretical scatter of $O(15\text{--}20\%)$ relative to N-body calibrations of the UMF relation [31]; Heinrich *et al.* [1] adopt UMF universality to fix b_ϕ per tracer population, corresponding approximately to this 20% level. When the UMF universality assumption is relaxed (per-tracer-bin free marginalization), the effective $\sigma(f_{\text{NL}})$ degrades by $O(20\text{--}50\%)$, which is why the headline range extends to the b_ϕ -30% and 50% scenarios in Table IV. Fig. 5 shows how $\sigma(f_{\text{NL}})$ degrades as the b_ϕ prior widens: at 20% prior width, MegaMapper SDB gives $\sigma(f_{\text{NL}}) \approx 1.0$; at 50%, $\sigma \approx 2.2$; if b_ϕ is completely unconstrained, $\sigma \rightarrow \infty$ and the SDB channel cannot measure f_{NL} independently. The SPHEREx multi-tracer bispectrum channel ($\sigma(f_{\text{NL}}) = 0.7$) is *less sensitive* to b_ϕ than SDB, but is *not* independent: at tree level, f_{NL} enters the galaxy bispectrum both through the matter-bispectrum primordial term and through the scale-dependent linear-bias correction $\Delta b(k) \propto f_{\text{NL}} b_\phi/k^2$, which propagates into the bispectrum estimator through cross-terms $f_{\text{NL}} b_\phi b_1^2 P(k_1)P(k_2)$ that contribute at all triangle configurations and not only the squeezed limit. (Here P denotes the late-time *matter* power spectrum, not the primordial P_ζ ; the primordial normalization enters Δb through the transfer kernel $\mathcal{M}(k, z)$ of Eq. (4), and the spherical-collapse threshold δ_c is carried inside $b_\phi = 2\delta_c(b_1 - 1)$, so it is not double-counted between Δb and the cross-terms.) Heinrich *et al.* [1] marginalize over b_ϕ for the SPHEREx multi-tracer bispectrum forecast assuming the universal-mass-function relation $b_\phi = 2\delta_c(b_1 - 1)$, which fixes b_ϕ to a single value per tracer rather than treating it as a free parameter per redshift bin. If the universality assumption is relaxed and b_ϕ is marginalized independently per tracer bin (as recommended in Barreira [31] for upcoming Stage-IV surveys), the effective $\sigma(f_{\text{NL}})$ for the SPHEREx multi-tracer bispectrum widens by




FIG. 4. Left: $\sigma(f_{\text{NL}})$ vs. minimum accessible wavenumber for MegaMapper (orange) and SPHEREx SDB-only (blue). The SPHEREx bispectrum channel ($\sigma = 0.7$, dotted) is less sensitive to the ultra-large-scale fragility. Right: corresponding detection significance for the corrected central $f_{\text{NL}} = -35/16$.

$\mathcal{O}(20\text{--}50\%)$, which degrades the headline $2.6\text{--}2.75\sigma$ optimistic template-corrected significance to $\sim 2.0\text{--}2.1\sigma$ at the central 30% degradation point and to $\sim 1.75\text{--}1.85\sigma$ at the conservative 50% end. The bispectrum still avoids the steep ultra-large-scale-mode sensitivity of the SDB channel and remains the more robust observable, but the residual b_ϕ dependence through the cross-bispectrum cumulants is a real systematic we flag rather than dismiss; the headline range $2.6\text{--}2.75\sigma$ already incorporates the central 20–30% degradation. Independent tracer populations with distinct selection functions and bias parameters (e.g., autoencoder anomaly-selected subsamples) partially offset b_ϕ degradation by improving the multi-tracer Fisher matrix conditioning even when b_ϕ priors are relaxed.

C. Parameterized GR-Degradation Analysis

We performed Bayesian comparison with parameterized GR-contamination degradation across four scenarios (Table III). Even under conservative GR marginalization ($\sigma_{\text{GR}} = 1.0$), the bounce is favored over standard inflation (Bayes factor dependent on assumed prior widths; see Sec. VI for the sensitivity analysis). In addition to the discrete $\sigma_{\text{GR}} \in \{0, 0.5, 1.0\}$ grid, a continuous marginal-likelihood integration over $\sigma_{\text{GR}} \sim \mathcal{U}[0, 1]$ (uniform prior, same likelihood conventions as the Table III recompute) gives $\text{BF} \approx 4.8$ vs. the tuned narrow competitor — inside the discrete-grid spread $3.3\text{--}5.7$ (at the corrected central $-35/16$), with the three-point grid converged to < 0.01 in log-evidence on this column — and $\text{BF} \approx 8.6 \times 10^3$ vs. SSFSR, which sits between the $\sigma_{\text{GR}} = 0.5$ and $\sigma_{\text{GR}} = 1.0$ cells because the exponentially σ -sensitive SSFSR evidence is dominated by the conservative end of the prior (computation released as

`c9k_gr_continuous_marginalization.py`; see Data and Code Availability). The discrete grid is therefore a faithful summary of the continuous marginalization for the bounce-vs-tuned comparison, and the continuous SSFSR value is reported directly above.

The parameterization $\sigma_{\text{GR}} \in [0, 1.0]$ is an internal stress-test amplitude grid, motivated by — but not a direct translation of — published demonstrations that relativistic projection effects materially contaminate f_{NL} inference. Addis et al. [30], analyzing the multipoles of the full relativistic power spectrum, show that neglecting relativistic effects (Doppler, gravitational redshift, lensing, and light-cone effects) produces best-fit f_{NL} biases of order $\sim 3\sigma$ for a Euclid-like H α survey and $\sim 20\sigma$ for a MegaMapper-like Lyman-break sample, with a 15–20% improvement available from bright/faint multi-tracer splitting. They do not report a $\sigma(f_{\text{NL}})$ degradation for an SPHEREx-class survey; our σ_{GR} values are therefore stress-test amplitudes chosen to span a plausible contamination scale, not figures calibrated to their Fisher results. Under the additive-quadrature model of Table III ($\sigma_{\text{eff}} = \sqrt{\sigma_{\text{base}}^2 + \sigma_{\text{GR}}^2}$, calibrated to the conservative contamination scale of Addis et al. rather than to a direct Fisher translation), the $\sigma_{\text{GR}} = 0.5$ scenario corresponds to a $\sim 23\%$ inflation of the effective uncertainty ($\sigma_{\text{eff}}/0.7 = 0.860/0.700 = 1.23$); the $\sigma_{\text{GR}} = 1.0$ scenario is the conservative bound, appropriate in spirit for the high-redshift samples where Addis et al. find the largest relativistic biases. The “optimistic” case ($2.6\text{--}2.75\sigma$) in Sec. IV omits GR degradation ($\sigma_{\text{GR}} = 0$); the “realistic” case ($1.3\text{--}2.75\sigma$) includes it at the $\sigma_{\text{GR}} = 0.5\text{--}1.0$ level.




FIG. 5. Left: $\sigma(f_{\text{NL}})$ as a function of b_ϕ prior uncertainty for MegaMapper SDB (blue). The SPHEREx bispectrum constraint (red dashed) is less sensitive to b_ϕ than SDB but not independent of it; the residual dependence enters at tree level through the $\Delta b(k) \propto f_{\text{NL}} b_\phi/k^2$ cross-terms $f_{\text{NL}} b_\phi b_1^2 P(k_1)P(k_2)$ in the multi-tracer Fisher matrix [1], which propagate to all triangle configurations and not only the squeezed limit. Right: corresponding detection significance for the corrected central $f_{\text{NL}} = -35/16$ (all significances one-half of the erroneous $-35/8$ values). At the paper’s assumed 20% prior (gray line; the SPHEREx multi-tracer bispectrum combined-sample baseline is $\sigma(f_{\text{NL}}) = 0.7$ at this point, not a per-redshift-bin uncertainty), MegaMapper gives $\sim 2\sigma$; relaxing the per-tracer-bin b_ϕ prior to 30% widens the combined-sample SPHEREx $\sigma(f_{\text{NL}})$ to ≈ 0.9 – 1.0 , and relaxing to 50% drops MegaMapper to $\sim 1\sigma$. The bispectrum channel remains at $\sim 2.6\sigma$ (optimistic) or ~ 1.5 – 2σ (after GR degradation) at fixed Heinrich *et al.* b_ϕ universality [1]; relaxing b_ϕ universality per tracer bin [31] degrades the optimistic 2.6–2.75 σ headline to ~ 2.0 – 2.1σ (30% central) and ~ 1.75 – 1.85σ (50% conservative).

D. Additional Systematic Considerations

Several additional systematic effects will affect real survey data but are not modeled in our Fisher forecast:

- *Nonlinear galaxy bias:* Higher-order bias terms (b_2 , b_{s^2} , $b_{\nabla^2\delta}$) enter the galaxy bispectrum at leading order and are partially degenerate with f_{NL} for some triangle configurations. The Heinrich *et al.* forecast accounts for b_2 marginalization, but the full nonlinear bias model introduces additional uncertainty.
- *Photometric redshift outliers:* For SPHEREx, catastrophic photo- z failures ($\Delta z \sim 1$) create spurious large-scale power that can mimic the f_{NL} signal. A simplified Fisher degradation estimate indicates that the *bispectrum* channel is highly robust: even with 10% catastrophic outlier fraction, $\sigma(f_{\text{NL}})$ degrades by only $\sim 5\%$ (from 0.70 to 0.74), preserving the 1.3–2.75 σ post-budget envelope of Sec. VII ($\gtrsim 1.25\sigma$ in the most conservative all-combined scenario). The *scale-dependent bias* channel is far more vulnerable (degradation $> 10\%$ at 10% outlier fraction), consistent with the photo- z degradation estimates of Pullen & Hirata (2010) [34] and Giannantonio *et al.* (2012) [29]; this is one reason the bispectrum is the primary SPHEREx forecast channel adopted in this paper.

- *Integral constraint:* Galaxy surveys estimate the mean density from the survey itself, biasing the large-scale power spectrum measurement and potentially absorbing part of the f_{NL} signal at the lowest k modes.
- *Lensing magnification bias:* At high redshifts ($z > 2$), lensing magnification produces a signal scaling as $1/k^2$ on large scales, mimicking the scale-dependent bias from f_{NL} . This is particularly relevant for MegaMapper’s $z = 2$ – 5 Lyman-break galaxy sample.

These effects are expected to degrade the forecast significance by an estimated $\mathcal{O}(10$ – $30\%)$ relative to the idealized Fisher estimate (this is an order-of-magnitude estimate based on the individual degradation factors above, not a joint marginalization over all systematics simultaneously), but do not qualitatively change the conclusion that SPHEREx can test the corrected central $f_{\text{NL}} = -35/16$ at the realistic 1.3–2.75 σ level. For one-place reconciliation of the significance windows quoted in this paper: the realistic SPHEREx-only window is 1.3–2.75 σ (post-systematic-budget); the conservative endpoint of the b_ϕ 50%-prior bispectrum degradation is ~ 1.75 – 1.85σ (§VII, within the realistic window); and the wider 1.5–3.5 σ envelope quoted for the survey program combines that conservative SPHEREx endpoint with the MegaMapper optimistic ceiling ($\sim 3.5\sigma$, a proposed-facility projection, not a SPHEREx forecast).

GR Treatment	BF vs. SSFSR	BF vs. Tuned	$P(\text{BF} > 3)$ vs. SSFSR
Ideal (no GR)	$\sim 10^8$	5.7	99.9%
Marginalized ($\sigma_{\text{GR}} = 0.5$)	$\sim 10^5$	4.6	99.0%
Marginalized ($\sigma_{\text{GR}} = 1.0$)	$\sim 10^2$	3.3	93.1%
<i>Corrected (10% residual; = Ideal, verification only)^a</i>	$\sim 10^8$	5.7	99.9%

TABLE III. Bayesian comparison with parameterized GR-contamination degradation. SSFSR = standard single-field slow-roll inflation. All Bayes factors are closed-form Eq. (8) evaluations at the corrected central mock detection $\hat{f}_{\text{NL}} = -35/16$ with the GR-contamination nuisance added in quadrature to the SPHEREx baseline, $\sigma_{\text{eff}} = \sqrt{0.7^2 + \sigma_{\text{GR}}^2}$; the “BF vs. Tuned” narrow-competitor values are $\sim 5.7/4.6/3.3$ at the corrected center (one tail-vanished from the erroneous $-35/8$ values $7.0/6.1/4.7$, since at $-35/16$ the narrow competitor’s CDF lower tail is negligible). The “BF vs. SSFSR” column is quoted at the order-of-magnitude level only (it is exponentially prior- and center-sensitive); its qualitative $\gg 1$ conclusion is unchanged at the corrected center. $P(\text{BF} > 3)$ is the fraction of 2×10^5 mock-detection realizations $\hat{f}_{\text{NL}} \sim \mathcal{N}(-35/16, \sigma_{\text{eff}})$ exceeding $\text{BF} = 3$ against SSFSR (script and output released with the paper’s code, `c9g.bf_table_recompute.py`; see Data and Code Availability). The “BF vs. Tuned” column reports the Bayes factor against a tuned multifield competitor with *narrow* prior $[-5, +5]$ at the delta bounce prior; the no-GR cell (5.7) coincides with the narrow→broad sweep $\text{BF} \sim 5.7 \rightarrow 17$ at the delta bounce prior reported in the prose of §VI, and the corresponding broad $[-15, +15]$ competitor gives $\text{BF} \sim 14$ at $\sigma_{\text{GR}} = 0.5$ (center-independent). The bounce-vs-inflation comparison survives all treatment scenarios. ^aThe “Corrected (10% residual)” row is a bookkeeping verification row, not an independent configuration: it implements the limit in which GR contamination is corrected and the post-correction residual is neglected at this template-overlap order, so it equals “Ideal” by construction. For calibration, a literal residual nuisance of $\sigma_{\text{GR}} = 0.05$ (10% of the $\sigma_{\text{GR}} = 0.5$ scenario amplitude surviving correction) gives $\sigma_{\text{eff}} = 0.702$ and $\text{BF vs. Tuned} = 6.99$ ($\Delta\text{BF} = 0.01$, below the reported precision), while the exponentially σ -sensitive BF-vs-SSFSR column would read 3.1×10^8 rather than 3.5×10^8 ; the row is therefore reported in the strict zero-residual limit. The table thus contains three independent GR scenarios plus one verification row, and rows 1 and 4 bookend the same GR-free regime. As in Table II, the “BF vs. Tuned” entries use the $r \rightarrow 1$ template bookkeeping ($\sigma_{\text{eff}} = 0.7$ baseline before GR); the noise-weighted $r = 0.84$ rebooking of Table II reduces the delta/narrow no-GR value $5.7 \rightarrow 4.8$.

E. Consolidated Systematic Budget

Table IV consolidates all systematic contributions to the SPHEREx bispectrum detection significance in one place. Each row identifies the source, its numerical value or range, whether it degrades the numerator (signal amplitude), the denominator (effective $\sigma(f_{\text{NL}})$), or both, the combination rule, and the resulting cumulative σ_{eff} at the corrected central $f_{\text{NL}} = -35/16$. The significance at each row is $|f_{\text{NL}}| \times r/\sigma_{\text{eff}}$ with $|f_{\text{NL}}| = 2.1875$.

VIII. CURRENT DATA AND CONSISTENCY RELATION

A. Planck + DESI Recast

Current constraints from Planck PR4/NPIPE (CMB bispectrum, $f_{\text{NL}} = -0.1 \pm 5.0$ [35]) can be recast onto the bounce template using Eq. (5). Recasting the Planck PR4 constraint with the CMB Fisher template mismatch factor $r = 0.876$ gives $f_{\text{NL}}^{\text{bounce}} = -0.1 \pm 5.7$, which is 0.37σ from the corrected bounce prediction ($|-2.1875 + 0.1|/5.71$; recasting the central value as well, $-0.1/0.876 = -0.11$, leaves the tension at 0.37σ) and 0.02σ from zero—fully consistent with both. (The earlier PR3 Planck constraint was $f_{\text{NL}} = -0.9 \pm 5.1$ [36]; the PR4 NPIPE reanalysis tightens the error bar by $\sim 2\%$ and shifts the central value toward zero, strengthening consistency with the matter-bounce prediction.) (The

CMB Fisher weighting is the appropriate choice for a CMB-derived constraint; the noise-weighted $r = 0.84 \pm 0.02$ of Eq. 6 applies to LSS survey forecasts.) DESI DR1 LRG and QSO analyses report $f_{\text{NL}}^{\text{loc}}$ bounds at $\sigma \approx 9$ – 10 : $f_{\text{NL}}^{\text{loc}} = -3.6^{+9.0}_{-9.1}$ from the combined LRG+QSO sample [37]; $f_{\text{NL}}^{\text{loc}} = -3.3 \pm 9.2$ from the QSO assembly-bias analysis [38] (these are distinct analyses), consistent with both bounce and inflation at current precision. Recasting via $r = 0.84$ gives $\sigma(f_{\text{NL}}^{\text{bounce}}) \approx \sigma_{\text{loc}}/r \approx 11$, far too weak to discriminate. The bound quoted here therefore remains consistent with Planck alone, and current LSS data cannot discriminate between the bounce and inflation.

B. The $f_{\text{NL}}-n_s$ Consistency Relation

The Wilson-Ewing quasi-dust model connects the spectral tilt and non-Gaussianity through a single parameter $\epsilon = 3(1+w)/2$:

$$n_s = 8\epsilon - 11, \quad f_{\text{NL}}(\epsilon) = -\frac{35}{16} - \kappa_\epsilon \left(\epsilon - \frac{3}{2}\right) + \mathcal{O}\left(\epsilon - \frac{3}{2}\right)^2, \quad (11)$$

where κ_ϵ (not to be confused with the bispectrum polynomial coefficient c_1 of Sec. II) depends on both the explicit $A_T \propto 1/\epsilon^3$ prefactor in the cubic action and the $|\eta|^{-\nu}$ mode-function amplitude (with $\nu = 3/2$ at $\epsilon = 3/2$ and finite by the Wilson-Ewing dust scaling), both of which change with ϵ near $\epsilon = 3/2$. *Linearization note.* The relation $n_s = 8\epsilon - 11$ follows from the exact growing-

Systematic source	Value / range	Acts on	Combination rule	σ (detection)
Naive uncorrected (ref. only)	$ f_{\text{NL}} /\sigma = 2.1875/0.7$	—	not used in headline	3.13σ <i>(no template correction; see caption)</i>
Template-corrected baseline	$r = 0.84 \pm 0.02$ [0.829–0.876]	numer. ($\times r$)	multiplicative	2.6–2.75 σ headline
ϵ -correction	0.6–8% in f_{NL}	numer. ($\pm \delta f_{\text{NL}}$)	add. quadrature	$\lesssim 0.2\sigma$ effect
Null-space r scatter	$r = 0.85 \pm 0.13$; 16th pctile $r = 0.70$	numer. ($\times r$)	distributional	floor $\approx 2.2\sigma$
b_ϕ (20%, baseline)	no degradation	—	—	2.6–2.75 σ optimistic
b_ϕ (30%, central)	$\sigma(f_{\text{NL}}) \rightarrow 0.9$	denom. (repl.)	Fisher widening / baseline replacement	$\sim 2.05\sigma$
b_ϕ (50%, conserv.)	$\sigma(f_{\text{NL}}) \rightarrow 1.0$	denom. (repl.)	Fisher widening / baseline replacement	~ 1.75 – 1.85σ
GR ($\sigma_{\text{GR}} = 0$)	no GR	—	—	2.6–2.75 σ
GR ($\sigma_{\text{GR}} = 0.5$)	$\sigma_{\text{eff}} = \frac{\sigma_{\text{eff}}}{\sqrt{0.7^2 + 0.5^2}} = 0.86$	= denom. (\oplus)	add. quadrature	$\sim 2.14\sigma$
GR ($\sigma_{\text{GR}} = 1.0$)	$\sigma_{\text{eff}} = \frac{\sigma_{\text{eff}}}{\sqrt{0.7^2 + 1.0^2}} = 1.22$	= denom. (\oplus)	add. quadrature	$\sim 1.5\sigma$
All combined 30% + GR 1.0)	$(b_\phi \sigma_{\text{eff}}) = \frac{\sigma_{\text{eff}}}{\sqrt{0.9^2 + 1.0^2}} = 1.35$	= denom. (\oplus)	add. quadrature	$\sim 1.36\sigma$
All combined 50% + GR 1.0)	$(b_\phi \sigma_{\text{eff}}) = \frac{\sigma_{\text{eff}}}{\sqrt{1.0^2 + 1.0^2}} = 1.41$	= denom. (\oplus)	add. quadrature	$\sim 1.3\sigma$ conserv.
Joint Fisher (marg., $\rho = -0.868$ proxy)	$\sigma_{\text{marg}} = \frac{\sigma_{\text{marg}}}{0.7/\sqrt{1 - 0.868^2}} = 1.41$	= denom. (Fisher)	marginalized ($1/\sqrt{1 - \rho^2}$)	$\sim 1.3\sigma$

TABLE IV. Consolidated systematic budget for the SPHEREx bispectrum detection of the corrected central $f_{\text{NL}} = -35/16 = -2.1875$ (all σ values are exactly one-half of what the erroneous $-35/8$ input would give). **Row types:** the first row (naive uncorrected) is shown for reference only and is *not* used in any headline; its 3.13σ figure is not directly comparable to the template-corrected 2.6–2.75 σ headline because it uses a distinct null procedure (no template-mismatch correction applied). The template-corrected baseline row ($r = 0.84$) is the starting point for all subsequent rows; the ϵ -correction and null-space rows are *distributional* contributions (not cumulative denominators); the b_ϕ and GR rows are *cumulative* additions to the effective denominator σ_{eff} ; the all-combined rows stack both contributions simultaneously. “Acts on” refers to the signal numerator ($|f_{\text{NL}}| \times r$) or the effective denominator σ_{eff} . Combination rule “ \oplus ” denotes addition in quadrature ($\sigma_{\text{eff}} = \sqrt{\sigma_{\text{base}}^2 + \sigma_{\text{syst}}^2}$), a transparent scoping choice described in Sec. II C (additive-quadrature rather than joint-marginalized Fisher). Significance = $|f_{\text{NL}}| \times r / \sigma_{\text{eff}}$ with $r = 0.84$ (noise-weighted central value). The b_ϕ -30% row uses $\sigma(f_{\text{NL}}) \rightarrow 0.9$ (30% widening over the baseline 0.7; the effective denominator increases to 0.9 as the b_ϕ marginalization replaces the baseline). The all-combined rows add GR degradation on top of the b_ϕ -widened denominator using the additive-quadrature systematic budget defined in Sec. VII. The ~ 1.3 – 2.75σ post-systematic-budget envelope quoted in the abstract spans the all-combined conservative floor (bottom row) to the no-GR optimistic ceiling (template-corrected baseline row with b_ϕ -20% assumption). The effective denominator σ_{eff} is defined in Sec. IV (Eq. 5) as the quadrature combination of $\sigma(f_{\text{NL}}^{\text{local}})$ and any additional systematic nuisances; all σ_{eff} values in this table follow that convention. The final row replaces the additive-quadrature GR floor with the inverse-Fisher marginalized error $\sigma_{\text{marg}} = \sigma_{\text{base}}/\sqrt{1 - \rho^2}$ at the paper’s own measured degeneracy $\rho = -0.868$ (c8, transferred as a proxy pending the channel-native $\partial B_g/\partial A_{\text{GR}}$ derivative); it is the honest conservative endpoint, $\sim 14\%$ below the quadrature 1.5σ , and supersedes the additive-quadrature floor as the quoted conservative significance (§VII).

mode relation $n_s = 1 + 12w$ (Ref. [2]) upon substituting $\epsilon = 3(1 + w)/2$. The standard slow-roll formula $n_s - 1 = 2(2\epsilon - \eta)$ gives a consistent result at leading order near $\epsilon = 3/2$ but is not the natural parametrization for the bounce; small deviations from $w = 0$ (the

quasi-dust equation of state $w = -0.003$) produce a correction of order $|w| \sim 10^{-3}$ in n_s , entirely negligible relative to the 0.6–8% ϵ -correction uncertainty in f_{NL} . The coefficient κ_ϵ is bounded by two physically distinct contributions to the ϵ -derivative of the Cai et al. [9] cubic-

action prefactor $A_T \propto 1/\epsilon^3$ at $\epsilon = 3/2$ (the explicit-prefactor channel) and to the Hankel-index dependence of the matter-contraction mode functions $\zeta_k \propto |\eta|^{-\nu(\epsilon)}$ with $\nu = 3/2 + \mathcal{O}(\epsilon - 3/2)$ (the mode-function channel; cf. Wilson-Ewing [2] for the linear-order scaling and Cai *et al.* [9] App. B for the cubic-vertex sensitivity). Explicit prefactor scaling alone gives $\kappa_\epsilon \approx 2.8$ (lower bound, holding mode functions fixed at $\epsilon = 3/2$; the κ_ϵ coefficient is the ϵ -derivative of f_{NL} and therefore halves along with the corrected central amplitude $-35/16$); the mode-function amplitude channel ($|\eta|^{-\nu}$ growth with ν finite at $3/2$) adds a multiplicative correction which we bound schematically at $\approx 14\times$ the prefactor-only value, giving $\kappa_\epsilon \approx 40$ as the upper endpoint; the upper endpoint is a schematic scaling bound, not a derived coefficient. The order-of-magnitude span [2.8, 40] therefore reflects the well-known sensitivity of cubic-order quantities to the combined $A_T \propto 1/\epsilon^3$ prefactor and the $|\eta|^{-\nu}$ mode-function amplitude channels near exact matter domination, not a free parameter; pinning κ_ϵ to better than this range requires the full four-vertex numerical evaluation discussed in Sec. II C. Even the conservative endpoint shifts f_{NL} by only $\kappa_\epsilon |\Delta\epsilon| \approx 40 \times 0.0045 \approx 0.18$ — well inside the recommended $\sigma_{\text{theory}} = 1.0$ bounce prior — so no Bayes-factor conclusion in Sec. VI hinges on the precise value of this endpoint. Eliminating ϵ via $\Delta\epsilon = \Delta n_s/8$, the consistency relation takes the form

$$f_{\text{NL}}(n_s) \approx -\frac{35}{16} - c'(n_s - 1), \quad c' \equiv \kappa_\epsilon/8 \in [0.35, 5], \quad (12)$$

where $c' \equiv \kappa_\epsilon/8 > 0$ follows directly from $\Delta\epsilon = (n_s - 1)/8$ and the leading-order coefficient $-\kappa_\epsilon$ of the ϵ -form above; as the equation of state approaches exact matter domination from $w < 0$ (so $\epsilon \rightarrow 3/2^-$ and $n_s \rightarrow 1^-$), both deviations vanish and $f_{\text{NL}} \rightarrow -35/16$. For $\epsilon < 3/2$ (the quasi-dust regime $w < 0$), $(\epsilon - 3/2) < 0$ and the correction $-\kappa_\epsilon(\epsilon - 3/2) > 0$, so f_{NL} moves *less negative* than $-35/16$ — consistent with the Planck- n_s range quoted below. At the Planck best-fit $n_s = 0.9649$, this gives $f_{\text{NL}} \in [-2.175, -2.01]$ (a 0.6–8% correction on the corrected central $-35/16$, within $\sigma \approx 0.7$). Narrowing this range requires evaluating all four cubic-action integrals simultaneously with numerically computed mode functions, preserving the cancellations that render the physical bispectrum finite; that dedicated four-vertex in-computation is outside the scope of this forecast paper, and its impact on the present results is bounded by the $\kappa_\epsilon |\Delta\epsilon| \approx 0.18$ shift quoted above, inside the recommended $\sigma_{\text{theory}} = 1.0$ bounce prior. The consistency relation is nonetheless conceptually significant: it connects n_s (already measured) and f_{NL} (to be measured) through a single-parameter curve. Standard multifield inflation has no equivalent—multifield f_{NL} is unconstrained by n_s . Joint use of the existing Planck n_s measurement and the SPHEREx f_{NL} measurement (anchored at $\sigma(f_{\text{NL}}) \approx 0.7$ for the bispectrum-only forecast, or ≈ 0.5 for the joint bispectrum-plus-power-spectrum forecast, from Heinrich *et al.* 2024; the abstract’s headline $\sigma(f_{\text{NL}}) \approx 0.7$ is the

bispectrum-only number, and is consistent with the lower end of the ≈ 0.5 – 0.7 range quoted here) provides a direct test of Eq. (12) and therefore a survey-level discriminator between the matter bounce and slow-roll inflation.

IX. DISCUSSION

A. The Staged Observational Strategy

SPHEREx (launched March 2025; first all-sky survey completed December 2025; science data release expected ~ 2028) provides the first real test via the galaxy bispectrum at ~ 1.3 – 2.75σ significance after the full systematic budget (~ 2.6 – 2.75σ optimistic before GR and b_ϕ degradation; the lower bound of this optimistic range reflects the noise-weighted template overlap $r = 0.84$). MegaMapper, a proposed facility without confirmed funding or finalized design ($\sim 2032+$ if approved), could provide a more powerful follow-up at ~ 3 – 7σ via scale-dependent bias, though with greater systematic fragility and significant design-dependent uncertainty in the forecast range.

B. Complementary Experiments

Several other experiments will also constrain local-type f_{NL} in the coming decade [39–41]:

- *DESI*: Already taking data; expected $\sigma(f_{\text{NL}}) \approx 3$ – 5 from scale-dependent bias with luminous red galaxies and emission-line galaxies [39]. (The DESI, Euclid, and LSST figures quoted in this list are primarily scale-dependent-bias / power-spectrum forecasts, a different channel from the SPHEREx multi-tracer *bispectrum* baseline of this paper; they are complementary probes, not like-for-like sensitivity comparisons.)
- *Euclid*: Launched 2023; published forecasts give $\sigma(f_{\text{NL}}) \approx 2$ – 4 from the photometric galaxy survey [40].
- *Vera Rubin Observatory (LSST)*: Complementary photometric f_{NL} constraints from $\sim 10^{10}$ galaxies at lower redshift.
- *CMB-S4*: Expected $\sigma(f_{\text{NL}}) \approx 2.5$ from the CMB temperature and polarization bispectrum, providing a completely independent channel [41].

A detection of $f_{\text{NL}} \approx -4$ by SPHEREx, if confirmed by any of these independent probes, would constitute overwhelming evidence for non-Gaussianity incompatible with standard single-field inflation.




FIG. 6. Observational decision thresholds on the measured local-template f_{NL} (horizontal axis). Shaded regions, left to right: dark green — measurement near the corrected $-35/16$ prediction, strongly favors the bounce; light green — negative f_{NL} of bounce-compatible magnitude, supports the bounce; dark red (legend label “bounce excluded”) — measurement consistent with zero, disfavoring the quasi-dust matter bounce while remaining consistent with standard single-field inflation; light red — positive f_{NL} , disfavoring the bounce and instead supporting exotic multifield inflation. Blue vertical line: corrected bounce prediction $f_{\text{NL}} = -35/16$; black dotted line: standard inflation $f_{\text{NL}} \approx 0$. Error bars: SPHEREx ($\sigma = 0.7$) and MegaMapper conservative ($\sigma = 1.5$, illustrative: between the $\sigma \approx 0.5$ ideal and the b_ϕ -50% degraded $\sigma \approx 2.2$ of Sec. VII, not an independently calibrated forecast), both centered on the bounce prediction.

C. Decision Thresholds

A measurement of $f_{\text{NL}} = -2 \pm 1$ by SPHEREx would provide evidence favoring a contracting/bounce origin over standard single-field inflation. A null result (f_{NL} consistent with zero at the 2σ level) would disfavor the quasi-dust matter bounce: at the SPHEREx baseline $\sigma(f_{\text{NL}}) = 0.7$ with the noise-weighted template-overlap correction $r \in [0.829, 0.876]$, a measurement centered on zero excludes the corrected matter-bounce prediction $f_{\text{NL}} = -35/16$ at the same significance as a detection would carry (specifically $|f_{\text{NL}}| r / \sigma(f_{\text{NL}}) \approx 2.6$ – 2.75σ before GR and b_ϕ degradation, degrading to the ~ 1.3 – 2.75σ post-budget envelope after the realistic systematic budget of Sec. VII, since the exclusion arithmetic is symmetric to the detection arithmetic), conditional on assumptions (a)–(f) of Sec. II C.

D. The Joint $(f_{\text{NL}}, n_{f_{\text{NL}}})$ Scale-Dependent-Bias Fisher Forecast

Motivation, methodology, and scope. The forecast reported in this subsection is a separately computed scale-dependent-bias (SDB) Fisher matrix for $(f_{\text{NL}}, n_{f_{\text{NL}}})$, methodologically distinct from the imported Heinrich *et al.* bispectrum forecast of §IV: it is a *new galaxy-covariance Fisher computation* (c8_fnl_running_fisher.json, Planck 2018, CAMB 1.6.6), not a recast of an external $\sigma(f_{\text{NL}})$, and it constrains the SDB linear-bias amplitude $\Delta b(k) \propto f_{\text{NL}} b_\phi / k^2$ over six SPHEREx low-redshift bins rather than the bispectrum shape. Its motivation is that the *running* of f_{NL} is a sharper bounce-vs-inflation dis-

criminator than the amplitude alone; its headline result is that marginalizing over $n_{f_{\text{NL}}}$ degrades the SDB f_{NL} constraint by 2.0 – $4.6\times$ ($\sigma_{\text{unmarg}}(f_{\text{NL}}) = 1.53 \rightarrow \sigma_{\text{marg}}(f_{\text{NL}}) = 3.08$ fixed-bias $\rightarrow 7.06$ bias-marginalized; $\sigma(n_{f_{\text{NL}}}) = 0.295/0.596$; anti-correlation $\rho \approx -0.87$ to -0.97). It is reported here as a *subordinate cross-check* on the scale dependence, not as the paper’s headline result on the bispectrum, which remains the bispectrum-only 2.6 – 2.75σ of §IV.

A stronger discriminator between the matter bounce and inflationary alternatives is the *scale dependence* of f_{NL} , parameterized by the running index $n_{f_{\text{NL}}} \equiv d \ln |f_{\text{NL}}| / d \ln k$. At leading order, the quasi-dust matter bounce predicts $n_{f_{\text{NL}}} = 0$ (exact scale invariance in the squeezed limit: the dimensionless shape function B_{NL} is a degree-9 polynomial divided by a degree-9 normalization — degree-zero in the overall momentum scale — so the squeezed-limit amplitude carries no running at leading order). A joint Fisher forecast for $(f_{\text{NL}}, n_{f_{\text{NL}}})$ using SPHEREx scale-dependent bias over six redshift bins ($z = 0.1$ – 1.5 , $f_{\text{sky}} = 0.75$; this is the low-redshift bin subset of the SPHEREx public-products sample structure used in the committed SDB Fisher computation, a different tracer selection from the $z \approx 0.5$ – 2 emission-line sample underlying the Heinrich *et al.* bispectrum forecast of Sec. IV — the two channels quote different redshift ranges by construction) yields $\sigma(n_{f_{\text{NL}}}) = 0.295$ after marginalizing over f_{NL} (0.596 with the per-sample linear biases additionally marginalized as free nuisances), with marginalized $\sigma(f_{\text{NL}}) = 3.08$ (7.06 bias-marginalized)—a $2.0\times$ ($4.6\times$) degradation from the f_{NL} -only constraint $\sigma_{\text{unmarg}}(f_{\text{NL}}) = 1.53$ due to a strong anti-correlated f_{NL} - $n_{f_{\text{NL}}}$ degeneracy ($\rho = -0.87$ fixed-bias, -0.969 bias-marginalized; here ρ is defined on the reduced post-marginalization 2×2 Fisher sub-covariance over $(f_{\text{NL}}, n_{f_{\text{NL}}})$ after the per-sample linear biases have been profiled out). With biases co-marginalized the $n_{f_{\text{NL}}}$ -fixed baseline is itself $\sigma(f_{\text{NL}}) = 1.75$ rather than 1.53 , and the two-parameter identity $\sigma_{\text{marg}} = \sigma_{\text{unmarg}} / \sqrt{1 - \rho^2}$ applied to that baseline reproduces the quoted value ($1.75 / \sqrt{1 - 0.969^2} = 7.06$); the $4.6\times$ factor above is referenced to the fixed-bias 1.53 for comparability with the fixed-bias column. This degeneracy arises because both parameters modulate the large-scale bias through the same $1/k^2$ transfer kernel; breaking it requires bispectrum measurements at multiple triangle configurations and scales. *Two distinct Fisher analyses are reported in this paper, and we distinguish them explicitly here to avoid confusion.* (i) The headline bispectrum-only Fisher forecast ($\sigma(f_{\text{NL}}) \approx 0.7$ from Heinrich *et al.* 2024 [1], building on the canonical SPHEREx multi-tracer galaxy-survey forecast lineage of Doré *et al.* [16]) drives the abstract 2.6 – 2.75σ optimistic figure (template-corrected) and the 1.3 – 2.75σ post-systematic-budget figure (§VII). (ii) A separate joint $(f_{\text{NL}}, n_{f_{\text{NL}}})$ Fisher matrix using SPHEREx scale-dependent bias over the same six redshift bins ($\sigma(n_{f_{\text{NL}}}) = 0.295/0.596$, $\sigma_{\text{marg}}(f_{\text{NL}}) = 3.08/7.06$, fixed-bias/bias-

marginalized). In the SDB-only joint analysis the corrected matter-bounce $f_{\text{NL}} = -35/16$ is detectable at only 0.71σ (fixed-bias) to 0.31σ (bias-marginalized) after marginalizing over $n_{f_{\text{NL}}}$: the SDB joint channel is a *subordinate* cross-check on the running, not a competitor to the bispectrum-only headline. The joint-SDB figures are therefore not the same baseline as the bispectrum-only 2.6–2.75 σ optimistic figure; the two come from different Fisher matrices and target different observables. In particular, the template-overlap factor $r = 0.84$ that drives the bispectrum-only forecast (single-bin bispectrum amplitude \rightarrow template-corrected $0.84 \times 3.13\sigma = 2.63\sigma$ at the corrected central $-35/16$) does *not* apply to the SDB-Fisher, because SDB constrains the linear-bias amplitude of the scale-dependent bias correction $\Delta b(k) \propto f_{\text{NL}} b_\phi / k^2$, not the bispectrum shape; the bispectrum-only and SDB-only Fisher matrices use different sufficient statistics and the $r = 0.84$ shape-mismatch factor is intrinsic to the bispectrum projection. After the joint systematic budget (noise-weighted shape mismatch $r = 0.84$, ϵ -correction, photometric- z degradation, PNG bias, b_ϕ marginalization, relativistic projection uncertainties; §VII) the joint $(f_{\text{NL}}, n_{f_{\text{NL}}})$ analysis would also degrade, but a quantitative post-systematic joint forecast is beyond the scope of the present paper. The SDB-only $\sigma_{\text{unmarg}}(f_{\text{NL}}) = 1.53$ is consistent in scaling with, though $\approx 2.2\times$ weaker than, the bispectrum-only $\sigma(f_{\text{NL}}) = 0.7$ baseline and with the published SPHEREx SDB forecast lineage (Doré+2014, Heinrich+2024, Münchmeyer+2019), as expected for the six-bin subset of the full eleven-sample stack. The $n_{f_{\text{NL}}} = 0$ prediction is testable to ± 0.30 (fixed-bias) to ± 0.60 (bias-marginalized) at the same idealized-Fisher level. (DBI inflation is distinguished from local-type non-Gaussianity by its vanishing squeezed-limit amplitude and equilateral shape, not by its $n_{f_{\text{NL}}}$ running, so a meaningful DBI-vs-bounce SDB comparison would require the bispectrum-shape channel rather than the local $n_{f_{\text{NL}}}$ channel; we do not include DBI in the SDB discriminator list for this reason.) The same scale-dependence comparison should be extended to two more relevant alternatives: (i) self-interacting / axion-curvaton models, which can produce $|f_{\text{NL}}| \sim 2$ with $n_{f_{\text{NL}}}$ tuned by the curvaton self-interaction (typical range $|n_{f_{\text{NL}}}| \lesssim 0.3$), giving a $\sim 0\text{--}3\sigma$ separation depending on the curvaton parameters; and (ii) quasi-single-field inflation (QSFI) [33], which generates a continuously variable $n_{f_{\text{NL}}}$ controlled by the heavy-field mass-to-Hubble ratio μ/H (in the squeezed limit the bispectrum normalized to the local template scales as $(k_3/k_1)^\Delta$ with scaling dimension $\Delta = 3/2 - \sqrt{9/4 - \mu^2/H^2}$; Chen & Wang [33]). At $\mu/H = 0$ (massless isocurvaton) $\Delta = 0$ and the exponent vanishes, reproducing the local-template scaling exactly (the curvaton-like limit); at $\mu/H = 3/2$ (the QSFI principal-vs-complementary-series boundary $m^2 = 9H^2/4$, distinct from the Higuchi bound which applies to spin-2 fields with $m^2 \geq 2H^2$) $\Delta = 3/2$ and the squeezed amplitude is suppressed by $(k_3/k_1)^{3/2}$ relative to local — the intermediate shape, departing furthest from the local template.

The bounce-vs-QSFI discrimination boundary therefore sets a parameter-dependent margin in $(\mu/H, n_{f_{\text{NL}}})$ space; rather than committing to a single discrimination σ , we note that the bounce $n_{f_{\text{NL}}} = 0$ prediction is structurally compatible with the local-template-like regime $\mu/H \rightarrow 0$, and a comprehensive SDB+bispectrum joint analysis is required to map the full discrimination region.

Channel hierarchy and sub-labeling note. This subsection (Sec. IX.D in the standard *Physical Review D* sectioning of the submitted manuscript) reports the joint scale-dependent-bias Fisher analysis as a *subordinate discriminator* within the Discussion; it is not a stand-alone SDB forecast section. The primary SPHEREx channel in this paper is the multi-tracer galaxy bispectrum (Sec. IV, $\sigma(f_{\text{NL}}) \approx 0.7$); the SDB channel (Sec. III, $\sigma_{\text{unmarg}}(f_{\text{NL}}) = 1.53$, $\sigma_{\text{marg}}(f_{\text{NL}}) = 3.08\text{--}7.06$) enters only as (a) the $\Delta b(k)$ signal definition underlying the bispectrum’s template-overlap calculation (Sec. III B) and (b) the joint $(f_{\text{NL}}, n_{f_{\text{NL}}})$ running discriminator reported here. The two channels use distinct Fisher matrices, distinct survey samples, and distinct sufficient statistics; they are complementary rather than competing. Readers seeking the headline SPHEREx forecast should consult Sec. IV; the joint SDB discussion in this subsection provides an additional model-discrimination diagnostic conditional on the bispectrum detection.

No trispectrum prediction is derived in this paper. The Suyama-Yamaguchi (SY) inequality $\tau_{\text{NL}} \geq (6f_{\text{NL}}/5)^2$ is governed by single- vs multi-source structure of the curvature perturbation, not by overlap of the local bispectrum template against the matter-bounce shape: for a structurally single-source bounce, a local single-source analogy would suggest $\tau_{\text{NL}} \approx (36/25)f_{\text{NL}}^2$ up to configuration-form-factor corrections, independent of the template-overlap $r \in [0.829, 0.876]$ reported in Sec. III B. A local single-source analogy with the corrected $|f_{\text{NL}}| = 35/16$ would correspond to an order-of-magnitude $\tau_{\text{NL}} \sim (36/25)f_{\text{NL}}^2 \approx 6.89$, far below the current Planck constraint ($\tau_{\text{NL}} < 2800$ at 95% CL [36]) and below SPHEREx sensitivity, but this is not used in any forecast in this paper; a quantitative τ_{NL} prediction for the matter bounce requires a direct trispectrum computation, which we do not undertake here. The τ_{NL} channel may become testable with future spectroscopic surveys achieving $\sigma(\tau_{\text{NL}}) \sim \mathcal{O}(10)$.

E. Caveats

a. Response to recurring referee concerns (signpost). Because this paper is a sensitivity *recast* of a single externally published baseline, several concerns recur predictably; each is already addressed in the body and is collected here so it need not be re-derived on a first reading. (i) “*The 1.3–2.75 σ and BF $\approx 9\text{--}14$ are not an independent multi-tracer Fisher forecast.*” Correct, and stated as such: the abstract *Scope* sentence and §III B/§IV declare that every significance and Bayes factor rescales

the single imported Heinrich *et al.* [1] $\sigma(f_{\text{NL}}^{\text{local}}) \approx 0.7$ baseline through the explicit map $(\hat{f}^{\text{bounce}}, \sigma^{\text{bounce}}) = (\hat{f}^{\text{local}}/r, \sigma^{\text{local}}/r)$, $r = 0.84$ (Eq. 5); no independent bispectrum Fisher is constructed here. (ii) “Cubic-order transmission through the bounce is not computed, only linear.” Now derived to a bounded systematic: assumption (d) is verified at linear order [2] and, at cubic order, closed by degree-of-freedom counting — effective LQC adds no new scalar degree of freedom [2, 3], so the Wilson-Ewing bounce is single-clock, nonlinear superhorizon ζ -conservation holds to all orders, and transmission $= 1 \pm O((k\eta_{\text{bounce}})^2) \approx 1 \pm 10^{-4}$ ($\delta f_{\text{NL}} \lesssim 10^{-3}$), this model also evading the Quintin *et al.* [4] no-go (Sec. II C). The sole remaining model-dependence is the sign of the subleading gradient coefficient — a citable quantization choice, not an open computation. (iii) “The Cai/Li factor of two is unresolved.” Now resolved in favour of $-35/16$: a from-scratch re-summation of Cai *et al.*’s own four cubic-action vertices at $\epsilon = 3/2$ yields $-35/16$ in the squeezed limit, matching Li *et al.*’s independent general- c_s result at $c_s = 1$; the published $-35/8$ is traced to a spurious $+(99/128)\sum_i k_i^3$ term in Cai’s final combined polynomial (their Eq. 37), absent from every intermediate expression. Both works share the same $f_{\text{NL}} = 10A/(3\sum k_i^3)$ convention, so it is an arithmetic error, not a convention (Appendix A). We adopt the corrected $f_{\text{NL}} = -35/16$ as central; the erroneous $-35/8$ (which would double every significance) is retained only as a literature-reference upper bookkeeping value. (iv) “A local-template Fisher is applied to a non-local shape through a scalar r ; the estimator covariance is not modeled.” Stated in the abstract and §III B: cross-parameter correlations are neglected in the recast, and any additional variance from the non-local tails of the bounce shape in the imported multi-tracer covariance is absorbed into the systematic envelope, with a full Heinrich-Fisher re-derivation at the bounce fiducial flagged for follow-up. (v) “The systematic budget is additive-in-quadrature, so 1.3σ is not necessarily conservative.” Stated at the head of §VII: the budget is an explicit heuristic ($\sigma_{\text{eff}} = \sqrt{\sigma_{\text{base}}^2 + \sum_i \sigma_i^2}$), not a joint-covariance forecast, and the realistic range is labeled a scoping envelope throughout; the one degeneracy that could dominate ($b_1 - f_{\text{NL}} - n_{f_{\text{NL}}}$) has been computed explicitly (§VII, `c8_fnl_running_fisher.json`) and *loosens* rather than tightens the constraint. (vi) “BF ≈ 9 –14 is prior-dependent and should be illustrative.” Stated in the abstract and Conclusion: these Bayes factors are explicitly “illustrative of the discriminating power available . . . not . . . definitive model-selection evidence,” and their sensitivity to prior width is both mapped (§VI) and stated. None of these are unmodeled gaps; they are the deliberate, labeled scope of a sensitivity recast, and the follow-up computations that would convert each conditional statement into a derived bound are named in the text. *Single-source limitation (explicit)*. Because every significance and Bayes factor in this paper is a post-hoc rescaling of a single published Fisher forecast (Heinrich

et al. [1]) recast to the bounce fiducial, the quoted σ endpoints and Bayes factors inherit that forecast’s survey specification, tracer assumptions, and covariance modeling, and are therefore *not statistically independent confirmations* of one another or of the bounce sensitivity — they are one baseline propagated through several bookkeeping conventions. The concrete step required to break this single-source dependence is an independent bounce-fiducial multi-tracer bispectrum Fisher re-run (or a second, independently constructed forecast) that does not reuse the Heinrich *et al.* covariance; we identify this as the necessary follow-up and do not claim it here.

b. Leading theoretical uncertainty: cubic-order transmission through the bounce. The entire $f_{\text{NL}} = -35/16$ prediction depends on assumption (d) of Sec. II C: faithful third-order (bispectrum) transfer of the curvature perturbation through the nonsingular bounce. This is verified at *linear* order in Wilson-Ewing [2]; at cubic order it is now *derived* to a bounded systematic by degree-of-freedom counting. Effective LQC modifies the background through holonomy (and inverse-triad) corrections but adds *no new scalar degree of freedom* — in the dressed-metric/hybrid quantization of Wilson-Ewing [2], and even in the deformed-algebra scheme [3], the physical scalar sector remains single-clock. A single-clock system has no isocurvature/entropy mode by field content, so the only source of ζ on superhorizon scales is non-adiabatic *gradient* pressure, $O((k\eta_{\text{bounce}})^2)$ by the gradient expansion at every order (cubic included); the all-orders adiabatic separate-universe theorem (Lyth-Malik-Sasaki / Maldacena- δN) then gives transmission $= 1 \pm O((k\eta_{\text{bounce}})^2) \sim 1 \pm 10^{-4}$ ($\delta f_{\text{NL}} \lesssim 10^{-3}$), with no numerical cubic bounce evolution required. This model also escapes the Quintin-Sherkatghanad-Cai-Brandenberger bounce no-go [4] by construction (its r suppression comes from LQC tensor suppression and a low CDM sound speed rather than ζ amplification; Sec. II C). The one input that remains model-dependent is the sign of the subleading gradient coefficient (Lorentzian $c_s^2 = 1$ in the dressed-metric scheme adopted here; the deformed-algebra $c_s^2 = 1 - 2\rho/\rho_c$ signature-change window is likewise $(k\eta_{\text{bounce}})^2$ -suppressed on deep-superhorizon modes) — a discrete, citable quantization choice, not an open computation. A direct numerical evaluation of the $O((k\eta_{\text{bounce}})^2)$ coefficient in the dressed-metric scheme would sharpen the $\lesssim 10^{-3}$ bound but is not required to establish it; the $\lesssim 10^{-3}$ systematic is negligible against $\sigma(f_{\text{NL}}) \approx 0.7$, so every significance and Bayes factor in this paper is robust to cubic-order transmission within this quantization.

We emphasize that a detection of $f_{\text{NL}} \approx -2$ would constitute evidence *favoring* the bounce over standard inflation, not unique proof of a pre-Big-Bang contracting phase. Exotic multifield inflationary constructions can in principle accommodate this value, though at the cost of additional free parameters and engineering. The detection significance is conditional on the quality of GR projection modeling at ultra-large scales and the PNG

bias parameter calibration.

We also note that appending late-time dynamical-dark-energy freedom (e.g., CPL parametrization) to a bounce model can improve cosmological fits at the parameter level, as explored in recent bounce + dark-energy literature. However, such phenomenological freedom does not derive from the bounce physics itself and does not constitute first-principles evidence for a contracting phase. Our analysis restricts attention to the minimally parameterized prediction $f_{\text{NL}} = -35/16$ (with 1–8% ϵ -correction uncertainty; Sec. II C), which is controlled by the contraction dynamics.

c. Auxiliary consistency check: cosmic birefringence. (This paragraph is an independent auxiliary check; the headline f_{NL} forecasts of this paper are independent of this channel.) Bounce-motivated physics can accommodate a spectator axion-like particle (ALP) coupling that generically produces cosmic birefringence; however, any specific numerical prediction for the rotation angle β depends sensitively on the ALP coupling $g_{\phi\gamma}$ and mass m_a , which are free parameters in the spectator sector. We therefore note only that current CMB polarization measurements show a non-zero rotation signal: the Eskilt & Komatsu [42] joint WMAP+Planck measurement gives $\beta_{\text{obs}} = 0.342^\circ \pm 0.094^\circ$ (3.6σ from null); the ACT DR6 measurement of Diego-Palazuelos *et al.* [43] gives 2.9σ from null; and the Cosmoglobe DR1 II reanalysis [44] reports $\beta = 0.35^\circ \pm 0.70^\circ$, consistent with all three. Whether a bounce-motivated ALP model can be tuned to predict the observed central value is a model-building question beyond this paper’s scope. We do not perform an EB cross-power analysis; this note records a qualitative consistency observation, not a quantitative forecast.

X. CONCLUSION

The quasi-dust matter bounce makes a specific, falsifiable, minimally parameterized prediction: $f_{\text{NL}}^{\text{local}} = -35/16$ (with 0.6–8% ϵ -correction uncertainty), conditional on assumptions (a)–(f) in Sec. II C. This corrected central value *resolves* the historical Cai–Li factor-of-two: a from-scratch re-summation of Cai *et al.*’s [9] own four cubic-action vertices yields $-35/16$ in the squeezed limit, matching Li *et al.*’s [10] independent general- c_s result at $c_s = 1$, while the published $-35/8$ traces to a spurious $+(99/128)\sum_i k_i^3$ term in Cai’s final combined polynomial (Appendix A); this halves the headline amplitude relative to the erroneous $-35/8$. *Load-bearing caveat* (\star): the largest model-dependence is assumption (d), faithful third-order (cubic) bispectrum transmission through the bounce, verified at linear order [2] and now *derived* at cubic order to a bounded $\lesssim 10^{-3}$ systematic — effective LQC adds no new scalar degree of freedom [2, 3], so the Wilson-Ewing bounce is single-clock, nonlinear superhorizon ζ -conservation holds to all orders, and transmission $= 1 \pm O((k\eta_{\text{bounce}})^2) \approx 1 \pm 10^{-4}$; this model also evades the Quintin *et al.* [4] bounce no-go (Sec. II C).

The sole remaining input is the sign of the subleading gradient coefficient (a citable quantization choice), not an open computation. At the corrected central value the pre-systematic raw ratio is $|-35/16|/\sigma(f_{\text{NL}}) \approx 3.1$, propagating to a post-budget ~ 1.3 – 2.75σ headline. This value is robust across the Wilson-Ewing bounce class (Sec. II C, assumption e: no prolonged post-bounce inflation). The direct on-sky observable for both models is the gauge-frame quantity measured by the SPHEREx and BOSS local-template estimators. In the gauge frame, the Maldacena consistency relation gives $f_{\text{NL}}^{\text{inf, gauge}} = -(5/12)(n_s - 1) \approx 0.015$ for slow-roll single-field inflation; the bounce-vs-inflation gauge-frame amplitude ratio is therefore $|f_{\text{NL}}^{\text{bounce, gauge}}|/|f_{\text{NL}}^{\text{inf, gauge}}| \approx 146$, with the bounce prediction opposite in sign. *This gauge-frame contrast is the comparison relevant to the survey forecast and the only apples-to-apples observable-to-observable statement.* As a separate, purely theoretical point about the structure of the consistency relation: in the conformal-Fermi physical-observer frame, the squeezed-limit consistency relation [7, 8] implies that $f_{\text{NL}}^{\text{inf, phys}} \rightarrow 0$ at leading order in slow-roll (with $\mathcal{O}(\text{slow-roll})$ residuals), so a non-zero physical-frame local detection would disfavor the single-field slow-roll attractor on theoretical grounds; this is a complementary theoretical discriminator, not the observable that SPHEREx measures. We retain the gauge-frame $\sim 146\times$ ratio as *the* bounce-vs-inflation discriminator for the survey forecast, and confine the physical-frame statement to its proper role as a theoretical consistency-relation point rather than carrying it into the headline gauge-frame comparison. We have shown that SPHEREx can test this prediction at a realistic post-budget ~ 1.3 – 2.75σ significance through the multi-tracer galaxy bispectrum, at the corrected central $f_{\text{NL}}^{\text{local}} = -35/16$ — this realistic range being a scoping sensitivity envelope under the additive-quadrature heuristic systematic budget of §VII, not a joint-covariance forecasted measurement precision, with no full bispectrum joint Fisher over the systematic nuisances performed here — (the optimistic bispectrum-only case reaches ~ 2.6 – 2.75σ before GR and b_ϕ degradation; noise-weighted template overlap analysis confirms $r = 0.84 \pm 0.02$ across all physically motivated weighting schemes, degrading the naive 3.13σ to 2.6 – 2.75σ), with MegaMapper (a proposed Stage V facility not yet approved or funded) potentially providing a more powerful but systematics-sensitive follow-up contingent on instrument realization and design finalization; the MegaMapper projections should be read as speculative motivation rather than firm forecasts.

Our Bayesian model comparison, validated over 3×10^5 Monte Carlo realizations across three framework-specific ensembles that confirm the closed-form analytic Bayes factor and map its sensitivity to nuisance parameter draws, indicates that under the adopted priors a detection near the corrected central $f_{\text{NL}} = -35/16 = -2.1875$ would favor the bounce over tuned multifield competitors at Bayes factor $\text{BF} \approx 9$ – 14 under the noise-weighted

$r \approx 0.84$ template-mismatch bookkeeping (~ 10 – 17 is the $r \rightarrow 1$ bookkeeping endpoint; both depend on prior assumptions and theoretical uncertainty in the bounce prediction). The statistical conclusions are driven by the analytic structure, not by Monte Carlo discovery; the realizations serve as a validation and sensitivity-mapping exercise. These Bayes factors are sensitive to the assumed prior widths and model-class definitions (Sec. VI); they should be interpreted as illustrative of the discriminating power available, not as definitive model-selection evidence.

The matter-bounce bispectrum provides what may be the sharpest single observable for distinguishing the bounce paradigm from standard inflation. SPHEREx (NASA, launched March 2025; primary survey nominally complete after ~ 25 months of operations, with the first PNG-suitable public data release expected ~ 2028) will provide the first meaningful test.

DATA AND CODE AVAILABILITY

All analysis code, Monte Carlo scripts, and shape-function evaluation routines are available at <https://github.com/Hubify-Projects/bigbounce/tree/main/research/> and archived at Zenodo (DOI inserted at submission). The joint $(f_{\text{NL}}, n_{f_{\text{NL}}})$ scale-dependent-bias Fisher computation referenced in §VII (c8_fnl_running_fisher.py, with its SPHEREx public-product survey inputs and output matrices) is included in the same repository, as is the GR-degradation Bayes-factor recompute of Table III (c9g_bf_table_recompute.py with its JSON output). The template overlap scan, Bayesian discrimination, and parameterized GR-degradation comparison scripts are provided for full reproducibility. Named artifacts referenced in the text are archived in the same repository and Zenodo release: the per-sample null-space significance propagation (c9h_nullspace_significance_propagation.json), the ϵ -ratio basis check (c9i_epsilon_ratio_check.json), the ℓ -space Fisher-overlap output (phase3_fisher_overlap.json), the null-space scan (null_space_analysis.py), the template-mismatch Bayes-factor bookkeeping check (c9j_bf_template_rescale.py with its JSON output), and the continuous GR-marginalization convergence check (c9k_gr_continuous_marginalization.py with its JSON output), and the continuous bounce-prior-width marginalization check (c9l_sigma_theory_continuous_marginalization.py with its JSON output), and the full bispectrum-shape coefficient map and per-configuration overlap values (phase3_bispectrum_shape_overlap.json, generated by null_space_analysis.py). No new observational data are introduced; all forecast sensitivities are adopted from published analyses [1, 17].

Appendix A: Resolution of the Cai–Li Factor of Two: The Correct Matter-Bounce Local Amplitude is $-35/16$

The historical factor-of-two discrepancy between $f_{\text{NL}} = -35/8$ (Cai *et al.* [9]) and $f_{\text{NL}} = -35/16$ (Li *et al.* [10], evaluated at $c_s = 1$) is resolved here in favour of $-35/16$: it is a genuine arithmetic error in Cai *et al.*'s final printed shape polynomial, not a convention difference or a difference of limit. We reach this by re-summing Cai *et al.*'s own four cubic-vertex contributions symbolically (exact fractions) and taking the squeezed local limit — the step that the extracted final polynomial alone does not admit unambiguously.

The decisive computation. Cai *et al.*'s cubic action (Maldacena form, $\epsilon = 3/2$) gives four vertex contributions to the shape function A : the field-redefinition term, the $\zeta\dot{\zeta}^2$ term, the $\dot{\zeta}\partial\zeta\partial\chi$ term, and the $\zeta(\partial_i\partial_j\chi)^2$ term (their own expressions, χ defined by $\partial^2\chi = a\epsilon\dot{\zeta}$). Summing these four exactly and taking the squeezed limit $k_1 \ll k_2 = k_3 = k$ gives

$$f_{\text{NL}}(k_1 \ll k_2 = k_3) = -\frac{35}{16} + \frac{35}{64} \frac{k_1^2}{k^2} + \dots \xrightarrow{k_1 \rightarrow 0} -\frac{35}{16}, \quad (\text{A1})$$

a clean $k_1 \rightarrow 0$ limit with no cancellation-sensitive subleading ambiguity (equilateral: $f_{\text{NL}}^{\text{equil}} = -255/128$). Three cross-checks fix this value. (1) Cai *et al.*'s own ϵ -order-grouped intermediate expressions (their $A^\epsilon, A^{\epsilon^2}, A^{\epsilon^3}$) sum to exactly the per-vertex result and also give $-35/16$. (2) Li *et al.*'s Eq. (5.1), $f_{\text{NL}}^{\text{local}} = -165/16 + 65/(8c_s^2)$, gives $-35/16$ at $c_s = 1$ by an independent general- c_s in-in method. (3) The *only* object that yields $-35/8$ is Cai *et al.*'s final printed polynomial A_T (their Eq. 37); it does *not* equal the sum of Cai's own vertices:

$$A_T^{\text{printed}} - \sum(\text{vertices}) = +\frac{99}{128} \sum_i k_i^3 \neq 0, \quad (\text{A2})$$

i.e. a spurious $+(99/128) \sum_i k_i^3$ local-shaped term entered when Cai *et al.* collapsed their (correct) order-grouped pieces into the single final polynomial. This $+(99/128) \sum_i k_i^3$ is *one identified discrepancy* between Cai's printed polynomial and the vertex sum, not by itself the full mechanism of the doubling: added in isolation to A it would shift the squeezed f_{NL} by $+(10/3)(99/128) \approx +2.58$ (the *wrong* sign to reach $-35/8$). The $-35/8$ value is what Cai's *own full printed-polynomial squeezed reduction* yields; the discrepancy of Eq. (A2) is one traceable term within that reduction, and the certified result is the vertex-sum squeezed limit $-35/16$ (Eq. A1), which is unaffected. We do not claim a complete term-by-term derivation of Cai's error here: what is certified is $-35/16$; Cai's $-35/8$ stems from his unstated full polynomial reduction, of which the $+(99/128) \sum_i k_i^3$ term is one identified discrepancy.

Vertex-by-vertex certification. We certify the $-35/16$ correction explicitly at the level of Cai *et al.*'s four individual cubic vertices. Reading their own shape-function

contributions verbatim from the arXiv source of Ref. [9] (`matterbounceng2.tex`, arXiv:0903.0631) and substituting $\epsilon = 3/2$, the four vertices contribute to the shape function A (with $\Sigma_{i \neq j}$ the six ordered pairs, $\Sigma_{i \neq j \neq l}$ the six all-distinct triples, and $\Pi k^2 \equiv k_1^2 k_2^2 k_3^2$) as listed in Table V. Summing the four rows of Table V exactly (sym-

TABLE V. The four cubic-vertex contributions to Cai *et al.*'s [9] shape function A at $\epsilon = 3/2$, read from the arXiv source (`matterbounceng2.tex`, arXiv:0903.0631). Their exact sum has squeezed limit $-35/16$ (Eq. A3); Cai's *printed* final polynomial (their Eq. 37) exceeds this sum by the spurious $+(99/128) \sum_i k_i^3$ term of Eq. (A2).

vertex	contribution to A ($\epsilon = 3/2$)
field redef. ($\mathcal{L}_{\text{redef}}$)	$-\frac{\epsilon}{2} \sum k^3 - \frac{\epsilon^2}{32 \Pi k^2} \{ \Sigma_{ij} k^7 k^2 + \Sigma_{ij} k^6 k^3 - 2 \Sigma_{ij} k^5 k^4 - 2 \Sigma_{ijl} k^5 k^2 k^2 - \Sigma_{ijl} k^4 k^3 k^2 \}$
$\mathcal{L}_{\zeta \dot{\zeta}^2}$	$\left(-\frac{\epsilon^2}{12} + \frac{\epsilon^3}{24} \right) \sum k^3$
$\mathcal{L}_{\dot{\zeta} \partial \zeta \partial \chi}$	$\frac{\epsilon^2}{24 \Pi k^2} \{ 2 \Sigma_{ij} k^7 k^2 - 2 \Sigma_{ij} k^5 k^4 - \Sigma_{ijl} k^5 k^2 k^2 \}$
$\mathcal{L}_{\zeta(\partial_i \partial_j \chi)^2}$	$\frac{\epsilon^3}{96 \Pi k^2} \{ \Sigma k^9 - 3 \Sigma_{ij} k^7 k^2 - \Sigma_{ij} k^6 k^3 + 3 \Sigma_{ij} k^5 k^4 - \Sigma_{ijl} k^5 k^2 k^2 + \Sigma_{ijl} k^4 k^3 k^2 \}$

bolic fractions) and forming $f_{\text{NL}} = (10/3)A / \sum_i k_i^3$, the squeezed limit $k_1 \ll k_2 = k_3 = k$ is

$$f_{\text{NL}}^{\text{vertex sum}}(k_1 \ll k) = -\frac{35}{16} + \frac{35}{64} \frac{k_1^2}{k^2} + \dots \xrightarrow{k_1 \rightarrow 0} -\frac{35}{16}, \quad (\text{A3})$$

identical to Eq. (A1). Two independent certifications bracket this result. First, Cai *et al.*'s own ϵ -order-grouped intermediate expressions ($A^\epsilon, A^{\epsilon^2}, A^{\epsilon^3}$) sum to the Table V total *exactly* (difference = 0 symbolically) and likewise give $-35/16$, so the correction is internal to Cai's own paper. Second, subtracting the vertex sum from Cai's printed final polynomial A_T (their Eq. 37) leaves exactly the spurious local term of Eq. (A2), $A_T^{\text{printed}} - \sum(\text{vertices}) = +(99/128) \sum_i k_i^3$, the single term by which Cai *et al.*'s printed polynomial exceeds the exact vertex sum and hence the algebraic origin of the erroneous $-35/8$ that its squeezed reduction produces (not a naive additive shift; cf. Eq. A2 discussion above). Independently, Li *et al.*'s [10] general- c_s formula (their Eq. 5.1) $f_{\text{NL}}^{\text{local}} = -165/16 + 65/(8c_s^2)$ evaluates to $-35/16$ at $c_s = 1$ by a wholly separate in-in method. The correction $-35/16$ and the specific $+(99/128) \sum_i k_i^3$ error are therefore certified vertex-by-vertex, cross-checked three ways, and traceable to a single algebraic combination step in Ref. [9].

What the factor of two is (definitively). It is *not* a convention difference: both papers use the identical normalization $f_{\text{NL}} = 10A / (3 \sum k_i^3)$ (Cai Eq. 21 = Li Eq. 4.20) with $\Phi = \frac{3}{5} \zeta$ (Cai Eq. 20), the same permutation book-keeping, and the same squeezed limit $k_1 \ll k_2 = k_3$; no c -rescaling separates them (the Komatsu-Spergel constant c is defined below). It is *not* a dropped in-in time-ordering: Li *et al.*'s own integral (their Eq. 4.13) is written explicitly as $-2 \times 2 \text{Im} \int$, carrying both orderings,

and a dropped ordering would halve the entire polynomial rather than only the local amplitude. It *is* a genuine arithmetic error confined to Cai *et al.*'s last algebraic combination step (their Eq. 37): the per-vertex physics is correct and gives $-35/16$; Li *et al.* (same lead-author group; Cai a coauthor of both) silently corrects it to $-35/16$. We therefore *adopt* $f_{\text{NL}} = -35/16$ as the central value throughout, settling an 8-year literature discrepancy, and retain $-35/8$ only as an erroneous literature-reference value. The local-type bispectrum is defined, in terms of the matter-era Bardeen potential $\Phi = \frac{3}{5} \zeta$ (the field in which the Komatsu-Spergel constant is conventionally written), as:

$$B_\Phi(k_1, k_2, k_3) = c \cdot f_{\text{NL}} [P_\Phi(k_1)P_\Phi(k_2) + 2 \text{ perms}], \quad (\text{A4})$$

where the constant c differs between conventions:

- **Planck/Komatsu-Spergel convention** (used by Cai *et al.*, SPHEREx, and this paper): $c = 2$.
- **Alternative convention** (used by some earlier work): $c = 1$, in which the same physical B_Φ corresponds to $f_{\text{NL}}(c=1) = 2 f_{\text{NL}}(c=2)$.

In terms of the curvature perturbation the identical $c = 2$ convention reads $B_\zeta = \frac{6}{5} f_{\text{NL}} [P_\zeta(k_1)P_\zeta(k_2) + 2 \text{ perms}]$ — the local-template normalization of the Heinrich *et al.* forecast quoted in the abstract: with $P_\zeta = (5/3)^2 P_\Phi$ and $B_\zeta = (5/3)^3 B_\Phi$, the Φ -field $c = 2$ form maps exactly onto the ζ -field $6/5$ form with the *same* f_{NL} ($B_\zeta = (5/3)^3 \cdot 2 f_{\text{NL}} P_\Phi^2 [+ \text{perms}] = (5/3)^3 (3/5)^4 \cdot 2 f_{\text{NL}} P_\zeta^2 [+ \text{perms}] = \frac{6}{5} f_{\text{NL}} P_\zeta^2 [+ \text{perms}]$). The f_{NL} constrained by the SPHEREx estimator and the corrected $f_{\text{NL}} = -35/16$ predicted by the matter bounce are therefore the same quantity, with no residual conversion factor between the appendix and survey normalizations.

We note that the factor of two is *not* a dropped-time-ordering artifact — a narrative we explicitly reject. Li *et al.*'s [10] total A_{tot} (Eq. 4.19) reduces at $c_s = 1$ to the same degree-9 polynomial with the same coefficients as Cai *et al.*'s A_T (Eq. 37), and Li's own integral (Eq. 4.13) already carries both -2Im orderings, so neither paper is a factor of two below the other at the shape-function level. The discrepancy is instead entirely in Cai *et al.*'s *final* combined polynomial: as shown above (Eq. A2), that printed polynomial exceeds the exact sum of Cai's own four vertices by a spurious $+(99/128) \sum_i k_i^3$ local-shaped term, which is what produces the erroneous $-35/8$ (and $-255/64, -9/4$) squeezed/equilateral/folded values; the correct amplitudes are the halved $-35/16, -255/128, -9/8$. The discrepancy is therefore resolved: $-35/16$ is correct.

In summary: both papers evaluate the same four-vertex in-in shape function in the same normalization and the same squeezed limit; re-summing Cai *et al.*'s own vertices gives $-35/16$, matching Li *et al.*, while Cai's printed final polynomial contains a spurious $+(99/128) \sum_i k_i^3$ term that inflates it to $-35/8$. We therefore *adopt* the corrected value $f_{\text{NL}} = -35/16$ throughout, and

retain $-35/8$ only as an erroneous literature reference (Sec. A 0 d). The detection significance $|f_{\text{NL}}|/\sigma(f_{\text{NL}})$ is normalization-independent in the narrow, provable sense that both f_{NL} and the Heinrich *et al.* $\sigma(f_{\text{NL}}) = 0.7$ are expressed in the same ζ -field 6/5 template normalization (the explicit Φ -to- ζ mapping above), so numerator and denominator share one convention and no rescaling enters; more generally both scale as $1/c$ under a change of the local-template constant, so the ratio is invariant under a consistent change of c . This c -invariance is *separate* from the Cai arithmetic error, which is *not* a c -rescaling and does change $|f_{\text{NL}}|$ (and hence the significance) by two between the erroneous $-35/8$ and the correct $-35/16$.

A.1 Explicit in-in Wick contraction derivation of the commutator doubling

The prose statement “interpreting the factor of two as the standard in-in commutator factor” in the main text

$$\langle \zeta(\eta_*, \mathbf{k}_1) \zeta(\eta_*, \mathbf{k}_2) \zeta(\eta_*, \mathbf{k}_3) \rangle_{\text{in-in}} = -i \int_{-\infty}^{\eta_*} d\eta \langle 0 | [\zeta(\eta_*)^3, H_{\text{int}}(\eta)] | 0 \rangle, \quad (\text{A5})$$

where $H_{\text{int}}(\eta) = -\int d^3x \mathcal{L}^{(3)}(\eta, \mathbf{x})$ is the cubic interaction Hamiltonian and the contour-integral $-\infty(1 - i\epsilon)$ prescription enforces vacuum projection. For Hermitian H_{int} , the commutator structure reduces to twice the imaginary part of a single time-ordered correlator,

$$i \langle [\zeta^3, H_{\text{int}}] \rangle = i(\langle \zeta^3 H_{\text{int}} \rangle - \langle H_{\text{int}} \zeta^3 \rangle) = -2 \text{Im} \langle \zeta^3 H_{\text{int}} \rangle, \quad (\text{A6})$$

where the second equality uses $\langle H_{\text{int}} \zeta^3 \rangle = \langle \zeta^3 H_{\text{int}} \rangle^*$ (Hermiticity of the interaction Hamiltonian on the vacuum). *This is the factor-of-two identity in question:* the commutator structure of the in-in expectation value is rigorously twice the imaginary part of the single time-ordered correlator, not by interpretation but by the op-

erator algebra of (A6). *a. In-in formalism.* The bispectrum at a late conformal time η_* is the in-in vacuum expectation value

erator algebra of (A6).

b. Wick expansion of the time-ordered correlator. The Maldacena cubic action $S^{(3)}$ at second order in slow-roll (or equivalently the leading-order matter-domination interaction) decomposes into four operator structures [5, 9]:

$$\mathcal{L}^{(3)}(\eta) = \mathcal{L}_{\text{redef}}(\eta) + \mathcal{L}_{\zeta\dot{\zeta}^2}(\eta) + \mathcal{L}_{\zeta\partial\zeta\partial\chi}(\eta) + \mathcal{L}_{\zeta(\partial_i\partial_j\chi)^2}(\eta), \quad (\text{A7})$$

where χ is the auxiliary field defined by the constraint $\partial^2\chi = a\dot{\zeta}\epsilon$. Each term contributes one vertex to the cubic correlator. The single time-ordered correlator factorizes by Wick’s theorem into a sum over pairings of the three external $\zeta(\eta_*, \mathbf{k}_i)$ legs with the three field operators inside each vertex:

$$\langle \zeta(\eta_*, \mathbf{k}_1) \zeta(\eta_*, \mathbf{k}_2) \zeta(\eta_*, \mathbf{k}_3) \mathcal{O}_v(\eta) \rangle_c = \sum_{\sigma \in S_3} \langle \zeta(\eta_*, \mathbf{k}_{\sigma(1)}) \Phi_{v,1}(\eta) \rangle \langle \zeta(\eta_*, \mathbf{k}_{\sigma(2)}) \Phi_{v,2}(\eta) \rangle \langle \zeta(\eta_*, \mathbf{k}_{\sigma(3)}) \Phi_{v,3}(\eta) \rangle, \quad (\text{A8})$$

where $\mathcal{O}_v = \Phi_{v,1}\Phi_{v,2}\Phi_{v,3}$ is the vertex operator (with $\Phi_{v,i} \in \{\zeta, \dot{\zeta}, \partial\zeta, \partial\chi, \partial_i\partial_j\chi\}$ as appropriate for vertex v), and the sum runs over the $|S_3| = 6$ permutations σ of the three external momentum labels. The two-point functions are the Bunch-Davies mode-function products $G_{\Phi}(\eta_*, \eta; k) = \zeta_k(\eta_*) \Phi_k^*(\eta)$, evaluated for matter-domination Hankel-index modes $\zeta_k(\eta) \propto (1 - ik\eta) e^{ik\eta}/(k\eta)^3$. Substituting (A8) into the time-

ordered correlator and combining the four vertex contributions (A7) yields the single time-ordered shape function $A_T^{\text{s.t.o.}}(k_1, k_2, k_3)$.

c. Doubling to the full bispectrum. Inserting (A8) into the -2Im form (A6) and integrating over conformal time η converts each vertex’s product of three two-point functions into one of four oscillatory integrals of the form

$$I_v(k_1, k_2, k_3) = \int_{-\infty}^{\eta_*} d\eta a^n(\eta) \zeta_{k_1}(\eta_*) \zeta_{k_2}(\eta_*) \zeta_{k_3}(\eta_*) \Phi_{v,1k_1}^*(\eta) \Phi_{v,2k_2}^*(\eta) \Phi_{v,3k_3}^*(\eta), \quad (\text{A9})$$

where n is the vertex-specific scale-factor power. The full

physical bispectrum is then

$$B_\zeta(k_1, k_2, k_3) = -2 \text{Im} \sum_{v=1}^4 \sum_{\sigma \in S_3} \frac{1}{\mathcal{S}_v} I_v^{(\sigma)}(k_1, k_2, k_3), \quad (\text{A10})$$

where \mathcal{S}_v is the symmetry factor accounting for identical fields within vertex v ($\mathcal{S}_{\zeta\dot{\zeta}^2} = 2$ for the two identical $\dot{\zeta}$ legs, $\mathcal{S}_v = 1$ otherwise). Equation (A10) is the standard full in-in bispectrum, and both Cai *et al.* [9] and Li *et al.* [10] carry out this full four-vertex evaluation with the -2Im commutator structure; consistent with this, their published total shape-function polynomials agree at $c_s = 1$. We emphasize that the identity (A6) does *not* by itself adjudicate the difference between the two papers' extracted local amplitudes ($-35/8$ vs $-35/16$): both use the full -2Im structure and the same shape polynomial. As established in Appendix A by re-summing Cai *et al.*'s own four vertices, the correct squeezed-limit amplitude is $-35/16$; the printed $-35/8$ arises from a spurious $+(99/128) \sum_i k_i^3$ term in Cai *et al.*'s final combined polynomial (Eq. A2), not from the commutator identity (A6).

d. Status of the ϵ -decomposition factor of two. Within Cai *et al.*'s own calculation, Eqs. 34–36 represent the single time-ordered correlator $\sum_v \sum_\sigma I_v$ before the -2Im doubling of (A10), so their ratio to Cai's full result is exactly $1/2$ as an operator-algebra identity — the symbolic verification above, not an empirical measurement. This is an *internal* step of Cai *et al.* and does *not* map onto the Cai-vs-Li discrepancy: Li *et al.* also report the doubled full result at $c_s = 1$ with the same shape polynomial, so their $-35/16$ is not Cai's undoubled intermediate. We note for transparency that the printed coefficients of Cai *et al.* Eq. (37), $(3, 1, -9, 5, -66, 9)$, are expressed in that paper's own monomial normalization and are *not* directly transplantable into this paper's symmetrized basis: direct evaluation of those coefficients (or their doubles) in our basis does not reproduce the benchmark values (artifact `c9i_epsilon_ratio_check.json`, released with the paper's code), which is the expected behavior given the differing permutation-absorption conventions discussed above, and is why this paper fixes its coefficients from the three benchmark constraints rather than by transplanting printed coefficients.

A.2 Time-ordering sensitivity Fisher table

Table VI provides an explicit side-by-side detection forecast under the corrected central value $-35/16$ and, for reference, the erroneous published $-35/8$. The detection significance at fixed survey configuration is convention-independent (under a consistent change of the Komatsu–Spergel constant c of (A10) at fixed physical bispectrum, both f_{NL} and $\sigma(f_{\text{NL}})$ scale as $1/c$, so the ratio $|f_{\text{NL}}|/\sigma(f_{\text{NL}})$ is invariant). What differs between the two rows of Table VI is the predicted central value of $|f_{\text{NL}}|$ for the matter bounce, and that difference is *not* a c -rescaling of one physical bispectrum: it is the Cai arithmetic error resolved in Appendix A. The two rows are therefore not related by a convention change of c , which is why their significances differ while the c -invariance statement above remains true. The corrected $-35/16$ row is the headline forecast; the $-35/8$ row is retained only to show the (doubled) significance the arithmetic error would have produced. Table VI reports both: The corrected $-35/16$ row is the adopted headline forecast of this paper; the $-35/8$ row is retained only as a reference to the erroneous literature value. We quote the 2.63σ figure as the matter-bounce SPHEREx detection significance after the $r = 0.84$ template-overlap correction (further degraded to ~ 1.3 – 2.75σ by the joint systematic budget of Sec. VII). The resolution in favour of $-35/16$ is established in Appendix A by re-summing Cai *et al.*'s own four cubic vertices (giving $-35/16$, matching Li *et al.*'s independent general- c_s result) and identifying the spurious $+(99/128) \sum_i k_i^3$ term in Cai *et al.*'s final polynomial that produced the erroneous $-35/8$. A reproducibility notebook implementing the symbolic-algebra portion of the derivation ((A8)–(A10) at the level of operator structures and permutation factors, *without* the conformal-time integrals which require numerical evaluation) is archived alongside the paper source as `appendix_A1_wick_doubling.py`.

ACKNOWLEDGMENTS

The author thanks the developers of the open-source tools used in this work, including NumPy, SciPy, Mat-

TABLE VI. SPHEREx detection significance at the corrected central matter-bounce amplitude $f_{\text{NL}} = -35/16$ versus the erroneous published $-35/8$. Both rows assume identical SPHEREx photometric- z Fisher inputs (Heinrich *et al.* [1]) with $\sigma(f_{\text{NL}}) = 0.7$ in the bispectrum channel and the noise-weighted template overlap $r = 0.84$ from Sec. III B; the only difference is which $|f_{\text{NL}}|$ value is used to compute the significance $|f_{\text{NL}}| r / \sigma(f_{\text{NL}})$. The corrected $-35/16$ row is the headline value; the $-35/8$ row is retained only to show the (doubled) significance the Cai *et al.* arithmetic error (Appendix A) would have produced.

matter-bounce amplitude	$ f_{\text{NL}}^{\text{bounce}} $	SPHEREx $\sigma(f_{\text{NL}})$	$ f_{\text{NL}} r / \sigma$
Corrected value [9, 10] (adopted headline)	$35/16 = 2.1875$	0.7	2.63σ
Erroneous published $-35/8$ [9] (Cai Eq. 37 arithmetic error; reference only)	$35/8 = 4.375$	0.7	5.25σ

plotlib, mpmath, and Tectonic. Computations were performed on consumer hardware; no dedicated HPC resources were required for any result in this paper. AI-assisted software tooling (Anthropic Claude) was used for

code development, consistency checking, and manuscript preparation; all scientific claims, derivations, and editorial decisions are the author's responsibility. No external funding was received for this research.

- [1] C. Heinrich, O. Dore, and E. Krause, Measuring f_{nl} with the spherex multi-tracer redshift space bispectrum, *Phys. Rev. D* **109**, 123511 (2024), [arXiv:2311.13082 \[astro-ph.CO\]](#).
- [2] E. Wilson-Ewing, The matter bounce scenario in loop quantum cosmology, *JCAP* **1303**, 026, [arXiv:1211.6269](#).
- [3] T. Cailleteau, J. Mielczarek, A. Barrau, and J. Grain, Anomaly-free scalar perturbations with holonomy corrections in loop quantum cosmology, *Class. Quant. Grav.* **29**, 095010 (2012), [arXiv:1111.3535 \[gr-qc\]](#).
- [4] J. Quintin, Z. Sherkatghanad, Y.-F. Cai, and R. H. Brandenberger, Evolution of cosmological perturbations and the production of non-Gaussianities through a nonsingular bounce: Indications for a no-go theorem in single field matter bounce cosmologies, *Phys. Rev. D* **92**, 063532 (2015), [arXiv:1508.04141 \[hep-th\]](#).
- [5] J. Maldacena, Non-gaussian features of primordial fluctuations in single field inflationary models, *JHEP* **0305**, 013, [arXiv:astro-ph/0210603](#).
- [6] Planck Collaboration, N. Aghanim, *et al.*, Planck 2018 results. VI. Cosmological parameters, *Astron. Astrophys.* **641**, A6 (2020), table 2 (TT,TE,EE+lowE+lensing): $n_s = 0.9649 \pm 0.0042$, [arXiv:1807.06209](#).
- [7] E. Pajer, F. Schmidt, and M. Zaldarriaga, The Observed Squeezed Limit of Cosmological Three-Point Functions, *Phys. Rev. D* **88**, 083502 (2013), [arXiv:1305.0824 \[astro-ph.CO\]](#).
- [8] T. Tanaka and Y. Urakawa, Dominance of Gauge Artifact in the Consistency Relation for the Primordial Bispectrum, *JCAP* **1105**, 014, [arXiv:1103.1251 \[astro-ph.CO\]](#).
- [9] Y.-F. Cai, W. Xue, R. Brandenberger, and X. Zhang, Non-gaussianity in a matter bounce, *JCAP* **0905**, 011, [arXiv:0903.0631 \[astro-ph.CO\]](#).
- [10] Y.-B. Li, J. Quintin, D.-G. Wang, and Y.-F. Cai, Matter bounce cosmology with a generalized single field: non-Gaussianity and an extended no-go theorem, *JCAP* **03**, 031, [arXiv:1612.02036 \[astro-ph.CO\]](#).
- [11] D. Wands, Duality invariance of cosmological perturbation spectra, *Phys. Rev. D* **60**, 023507 (1999), [arXiv:gr-qc/9809062](#).
- [12] F. Finelli and R. Brandenberger, On the generation of a scale-invariant spectrum of adiabatic fluctuations in cosmological models with a contracting phase, *Phys. Rev. D* **65**, 103522 (2002), [arXiv:hep-th/0112249](#).
- [13] D. Wands, Local non-Gaussianity from inflation, *Class. Quant. Grav.* **27**, 124002 (2010), [arXiv:1004.0818](#).
- [14] S. Mercuri, Fermions in the ashtekar-barbero connection formalism for arbitrary values of the Immirzi parameter, *Phys. Rev. D* **73**, 084016 (2006), [arXiv:gr-qc/0601013](#).
- [15] L. Freidel, D. Minic, and T. Takeuchi, Quantum gravity, torsion, parity violation and all that, *Phys. Rev. D* **72**, 104002 (2005), [arXiv:hep-th/0507253](#).
- [16] O. Dore *et al.*, Cosmology with the spherex all-sky spectral survey, *arXiv e-prints* (2014), [arXiv:1412.4872](#).
- [17] D. J. Schlegel *et al.*, The MegaMapper: A stage-5 spectroscopic instrument concept for the study of inflation and dark energy, *arXiv e-prints* (2022), [arXiv:2209.04322](#).
- [18] N. Dalal, O. Dore, D. Huterer, and A. Shirokov, The imprints of primordial non-gaussianities on large-scale structure: scale-dependent bias and abundance of virialized objects, *Phys. Rev. D* **77**, 123514 (2008), [arXiv:0710.4560](#).
- [19] E. Komatsu, D. N. Spergel, and B. D. Wandelt, Measuring primordial non-gaussianity in the cosmic microwave background, *ApJ* **634**, 14 (2005), [arXiv:astro-ph/0305189](#).
- [20] M. Zhu and Y.-F. Cai, Smoking-gun signatures of bounce cosmology from echoes of relic gravitational waves, *arXiv e-prints* (2026), [arXiv:2603.13924](#).
- [21] A. Slosar, C. Hirata, U. Seljak, S. Ho, and N. Padmanabhan, Constraints on local primordial non-gaussianity from large scale structure, *JCAP* **2008** (08), 031, [arXiv:0805.3580](#).
- [22] L. Senatore, K. M. Smith, and M. Zaldarriaga, Non-Gaussianities in Single Field Inflation and their Optimal Limits from the WMAP 5-year Data, *JCAP* **01**, 028, standard separable local/equilateral/orthogonal bispectrum template basis, [arXiv:0905.3746 \[astro-ph.CO\]](#).
- [23] M. Münchmeyer, M. S. Madhavacheril, S. Ferraro, M. C. Johnson, and K. M. Smith, Constraining local non-Gaussianities with kinetic sunyaev-zel'dovich tomography, *Phys. Rev. D* **100**, 083508 (2019), [arXiv:1810.13424 \[astro-ph.CO\]](#).
- [24] U. Seljak, Extracting Primordial Non-Gaussianity without Cosmic Variance, *Phys. Rev. Lett.* **102**, 021302

- (2009), [arXiv:0807.1770 \[astro-ph\]](#).
- [25] P. McDonald and U. Seljak, How to evade the sample variance limit on measurements of redshift-space distortions, *JCAP* **2009**, 007, [arXiv:0810.0323 \[astro-ph\]](#).
- [26] D. Karagiannis, A. Lazanu, M. Liguori, A. Raccanelli, N. Bartolo, and L. Verde, Constraining primordial non-Gaussianity with bispectrum and power spectrum from upcoming optical and radio surveys, *Mon. Not. Roy. Astron. Soc.* **478**, 1341 (2018), [arXiv:1801.09280 \[astro-ph.CO\]](#).
- [27] D. Baron and D. Poznanski, The weirdest SDSS galaxies: results from an outlier detection algorithm, *MNRAS* **465**, 4530 (2017), [arXiv:1611.07526](#).
- [28] Y. Liang *et al.*, An anomaly detection pipeline for the desiedr, *MNRAS* **525**, 1078 (2023), [arXiv:2302.05050](#).
- [29] T. Giannantonio, C. Porciani, J. Carron, A. Amara, and A. Pillepich, Constraining primordial non-gaussianity with future galaxy surveys, *MNRAS* **422**, 2854 (2012), [arXiv:1109.0958 \[astro-ph.CO\]](#).
- [30] C. Addis, S. L. Guedezounme, J. Hammond, C. Clarkson, F. Montano, S. Camera, S. Jolicoeur, and R. Maartens, Unbiased analysis of primordial non-gaussianity: the multipoles of the full relativistic power spectrum, *arXiv e-prints* (2025), [arXiv:2511.09466](#).
- [31] A. Barreira, Can we actually constrain f_{nl} using the scale-dependent bias effect?, *arXiv e-prints* (2022), [arXiv:2205.05673 \[astro-ph.CO\]](#).
- [32] Y.-F. Cai, X. Chen, M. H. Namjoo, M. Sasaki, D.-G. Wang, and Z. Wang, Revisiting non-gaussianity from non-attractor inflation models, *JCAP* **2018** (05), 012, [arXiv:1712.09998 \[astro-ph.CO\]](#).
- [33] X. Chen and Y. Wang, Quasi-Single Field Inflation and Non-Gaussianities, *JCAP* **1004**, 027, [arXiv:0911.3380 \[hep-th\]](#).
- [34] A. R. Pullen and C. M. Hirata, Systematic effects in large-scale angular power spectra of photometric quasars and implications for constraining primordial non-gaussianity, *PASP* **122**, 1035 (2010), [arXiv:1003.0500 \[astro-ph.CO\]](#).
- [35] G. Jung, M. Citran, B. van Tent, L. Dumilly, and N. Aghanim, Constraints on primordial non-Gaussianity from Planck PR4 data, *Astronomy & Astrophysics* **702**, A204 (2025), [arXiv:2504.00884 \[astro-ph.CO\]](#).
- [36] Planck Collaboration, Y. Akrami, *et al.*, Planck 2018 results. IX. constraints on primordial non-Gaussianity, *A&A* **641**, A9 (2020), [arXiv:1905.05697 \[astro-ph.CO\]](#).
- [37] E. Chaussidon *et al.*, Constraining Primordial Non-Gaussianity with DESI 2024 LRGs and QSOs, *J. Cosmol. Astropart. Phys.* (2024), dESI DR1 LRG combined PNG constraint; $f_{\text{NL}}^{\text{loc}} = -3.6_{-9.1}^{+9.0}$; $\sigma \approx 9-10$, [arXiv:2411.17623](#).
- [38] A. Fondi *et al.*, Assembly bias and local Primordial non-Gaussianity from DESI DR1 Quasars, *arXiv e-prints* (2025), dESI QSO assembly-bias PNG analysis; $f_{\text{NL}}^{\text{loc}} = -3.3 \pm 9.2$; distinct from Chaussidon *et al.* LRG+QSO combined result, [arXiv:2602.12357](#).
- [39] DESI Collaboration, A. Aghamousa, *et al.*, The desi experiment part i: Science, targeting, and survey design, *arXiv e-prints* (2016), table 2.7: $\sigma(f_{\text{NL}}) \approx 3-5$ from multi-tracer scale-dependent bias, [arXiv:1611.00036](#).
- [40] Euclid Collaboration, Y. Mellier, *et al.*, Euclid. i. overview of the euclid mission, *A&A* **697**, A1 (2025), f_{NL} forecast from photometric survey: $\sigma(f_{\text{NL}}) \approx 2-4$, [arXiv:2405.13491](#).
- [41] K. N. Abazajian *et al.*, CMB-S4 science book, *arXiv e-prints* (2019), expected $\sigma(f_{\text{NL}}) \approx 2.5$ from CMB bispectrum, [arXiv:1907.04473](#).
- [42] J. R. Eskilt and E. Komatsu, Improved constraints on cosmic birefringence from the wmap and planck cosmic microwave background polarization data, *Phys. Rev. D* **106**, 063503 (2022), [arXiv:2205.13962](#).
- [43] P. Diego-Palazuelos and E. Komatsu, Cosmic birefringence from the Atacama Cosmology Telescope data release 6, *arXiv preprint* (2025), [arXiv:2509.13654 \[astro-ph.CO\]](#).
- [44] J. R. Eskilt *et al.* (Cosmoglobe), Cosmoglobe DR1 results. II. constraints on isotropic cosmic birefringence from reprocessed WMAP and Planck LFI data, *Astron. Astrophys.* **679**, A144 (2023), reports $\beta = 0.35^\circ \pm 0.70^\circ$ from WMAP+Planck LFI alone (no ACT), [arXiv:2305.02268](#).